

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
5 July 2001 (05.07.2001)

PCT

(10) International Publication Number  
**WO 01/47959 A2**

(51) International Patent Classification<sup>7</sup>: C07K 14/00

(21) International Application Number: PCT/US00/42334

(22) International Filing Date:  
29 November 2000 (29.11.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/451,739 30 November 1999 (30.11.1999) US  
09/602,362 24 October 2000 (24.10.2000) US

(71) Applicants (for all designated States except US): LUDWIG INSTITUTE FOR CANCER RESEARCH [CH/US]; 605 Third Avenue, New York, NY 10158 (US). MEMORIAL SLOAN-KETTERING CANCER CENTER [US/US]; 1275 York Avenue, New York, NY 10021 (US). CORNELL RESEARCH FOUNDATION [US/US]; 20 Thornwood Drive, Ithaca, NY 11850 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): JAGER, Dirk [—/US]; 1275 York Avenue, New York, NY 10021 (US).

STOCKERT, Elisabeth [—/US]; 1275 York Avenue, New York, NY 10021 (US). SCANLAN, Matthew [—/US]; 1275 York Avenue, New York, NY 10021 (US). KNUTH, Alexander [—/DE]; Steinbacher Hohl 2-28, 60488 Frankfurt am Main (DE). OLD, Lloyd [—/US]; 605 Third Avenue, New York, NY 10158 (US). GURE, Ali [—/US]; 1275 York Avenue, New York, NY 10021 (US). CHEN, Yao-Tseng [—/US]; 525 East 68th Street, New York, NY 10021 (US).

(74) Agent: HANSON, Norman, D.; Fulbright & Jaworski L.L.P., 666 Fifth Avenue, New York, NY 10103 (US).

(81) Designated States (national): AU, CA, CN, JP, KR, US.

(84) Designated States (regional): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR).

**Published:**

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/47959 A2

(54) Title: ISOLATED NUCLEIC ACID MOLECULES ENCODING CANCER ASSOCIATED ANTIGENS, THE ANTIGENS PER SE, AND USES THEREOF

(57) Abstract: The invention relates to newly identified cancer associated antigens. It has been discovered that each of these molecules provokes antibodies when expressed by a subject. The ramifications of this observation are also a part of this invention.



ISOLATED NUCLEIC ACID MOLECULES ENCODING CANCER ASSOCIATED  
ANTIGENS, THE ANTIGENS PER SE, AND USES THEREOF

RELATED APPLICATIONS

This application is a continuation in part of Serial No. 09/602, 362, filed June 22, 2000 which is a continuation in part of Serial No. 09/451,739, filed November 30, 1999, both of which are incorporated by reference in their entirety.

FIELD OF THE INVENTION

This invention relates to antigens associated with cancer, the nucleic acid molecules encoding them, as well as the uses of these.

BACKGROUND AND PRIOR ART

It is fairly well established that many pathological conditions, such as infections, cancer, autoimmune disorders, etc., are characterized by the inappropriate expression of certain molecules. These molecules thus serve as "markers" for a particular pathological or abnormal condition. Apart from their use as diagnostic "targets", i.e., materials to be identified to diagnose these abnormal conditions, the molecules serve as reagents which can be used to generate diagnostic and/or therapeutic agents. A by no means limiting example of this is the use of cancer markers to produce antibodies specific to a particular marker. Yet another non-limiting example is the use of a peptide which complexes with an MHC molecule, to generate cytolytic T cells against abnormal cells.

Preparation of such materials, of course, presupposes a source of the reagents used to generate these. Purification from cells is one laborious, far from sure method of doing so.

Another preferred method is the isolation of nucleic acid molecules which encode a particular marker, followed by the use of the isolated encoding molecule to express the desired molecule.

Two basic strategies have been employed for the detection of such antigens, in e.g., human tumors. These will be referred to as the genetic approach and the biochemical approach. The genetic approach is exemplified by, e.g., dePlaen et al., Proc. Natl. Sci. USA 85: 2275 (1988), incorporated by reference. In this approach, several hundred pools of plasmids of a cDNA library obtained from a tumor are transfected into recipient cells, such as COS cells, or into antigen-negative variants of tumor cell lines which are tested for the expression of the specific antigen. The biochemical approach, exemplified by, e.g., O. Mandelboim, et al., Nature 369: 69 (1994) incorporated by reference, is based on acidic elution of peptides which have bound to MHC-class I molecules of tumor cells, followed by reversed-phase high performance liquid chromatography (HPLC). Antigenic peptides are identified after they bind to empty MHC-class I molecules of mutant cell lines, defective in antigen processing, and induce specific reactions with cytotoxic T-lymphocytes. These reactions include induction of CTL proliferation, TNF release, and lysis of target cells, measurable in an MTT assay, or a  $^{51}\text{Cr}$  release assay.

These two approaches to the molecular definition of antigens have the following disadvantages: first, they are enormously cumbersome, time-consuming and expensive; and second, they depend on the establishment of cytotoxic T cell lines (CTLs) with predefined specificity.

The problems inherent to the two known approaches for the identification and molecular definition of antigens is best demonstrated by the fact that both methods have, so far, succeeded in defining only very few new antigens in human tumors. See, e.g., van der Bruggen et al., Science 254: 1643-1647 (1991); Brichard et al., J. Exp. Med. 178: 489-495 (1993); Coulie, et

al., J. Exp. Med. 180: 35-42 (1994); Kawakami, et al., Proc. Natl. Acad. Sci. USA 91: 3515-3519 (1994).

Further, the methodologies described rely on the availability of established, permanent cell lines of the cancer type under consideration. It is very difficult to establish cell lines from certain cancer types, as is shown by, e.g., Oettgen, et al., Immunol. Allerg. Clin. North. Am. 10: 607-637 (1990). It is also known that some epithelial cell type cancers are poorly susceptible to CTLs in vitro, precluding routine analysis. These problems have stimulated the art to develop additional methodologies for identifying cancer associated antigens.

One key methodology is described by Sahin, et al., Proc. Natl. Acad. Sci. USA 92: 11810-11913 (1995), incorporated by reference. Also, see U.S. Patent No. 5,698,396, and Application Serial No. 08/479,328, filed on June 7, 1995 and January 3, 1996, respectively. All three of these references are incorporated by reference. To summarize, the method involves the expression of cDNA libraries in a prokaryotic host. (The libraries are secured from a tumor sample). The expressed libraries are then immunoscreened with absorbed and diluted sera, in order to detect those antigens which elicit high titer humoral responses. This methodology is known as the SEREX method ("Serological identification of antigens by Recombinant Expression Cloning"). The methodology has been employed to confirm expression of previously identified tumor associated antigens, as well as to detect new ones. See the above referenced patent applications and Sahin, et al., supra, as well as Crew, et al., EMBO J 144: 2333-2340 (1995).

This methodology has been applied to a range of tumor types, including those described by Sahin et al., supra, and Pfreundschuh, supra, as well as to esophageal cancer (Chen et al., Proc. Natl. Acad. Sci. USA 94: 1914-1918 (1997)); lung cancer (Güre et al., Cancer Res. 58:

1034-1041 (1998)); colon cancer (Serial No. 08/948, 705 filed October 10, 1997) incorporated by reference, and so forth. Among the antigens identified via SEREX are the SSX2 molecule (Sahin et al., Proc. Natl. Acad. Sci. USA 92: 11810-11813 (1995); Tureci et al., Cancer Res. 56: 4766-4772 (1996); NY-ESO-1 Chen, et al., Proc. Natl. Acad. Sci. USA 94: 1914-1918 (1997); and SCP1 (Serial No. 08/892,705 filed July 15, 1997) incorporated by reference. Analysis of SEREX identified antigens has shown overlap between SEREX defined and CTL defined antigens. Mage-1, tyrosinase, and NY-ESO-1 have all been shown to be recognized by patient antibodies as well as CTLs, showing that humoral and cell mediated responses do act in concert.

It is clear from this summary that identification of relevant antigens via SEREX is a desirable aim. The inventors have applied this methodology and have identified several new antigens associated with cancer, as detailed in the description which follows.

#### DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

##### **EXAMPLE 1**

The SEREX methodology, as described by, e.g. Sahin, et al., Proc. Natl. Acad. Sci. USA 92: 11810-11813 (1995); Chen, et al., Proc. Natl. Acad. Sci. USA 94: 1914-1918 (1997), and U.S. Patent No. 5,698,396, all of which are incorporated by reference. In brief, total RNA was extracted from a sample of a cutaneous metastasis of a breast cancer patient (referred to as "BR11" hereafter), using standard CsCl guanidine thiocyanate gradient methodologies. A cDNA library was then prepared, using commercially available kits designed for this purpose. Following the SEREX methodology referred to supra, this cDNA expression library was amplified, and screened with either autologous BR11 serum which had been diluted to 1:200, or with allogeneic, pooled serum, obtained from 7 different breast cancer patients, which had been

diluted to 1:1000. To carry out the screen, serum samples were first diluted to 1:10, and then preabsorbed with lysates of *E. coli* that had been transfected with naked vector, and the serum samples were then diluted to the levels described supra. The final dilutions were incubated overnight at room temperature with nitrocellulose membranes containing phage plaques, at a density of 4-5000 plaque forming units ("pfus") per 130 mm plate.

Nitrocellulose filters were washed, and incubated with alkaline phosphatase conjugated, goat anti-human Fc $\gamma$  secondary antibodies, and reactive phage plaques were visualized via incubation with 5-bromo-4-chloro-3-indolyl phosphate and nitroblue tetrazolium.

This procedure was also carried out on a normal testicular cDNA library, using a 1:200 serum dilution.

A total of  $1.12 \times 10^6$  pfus were screened in the breast cancer cDNA library, and 38 positive clones were identified. With respect to the testicular library,  $4 \times 10^5$  pfus were screened, and 28 positive clones were identified.

Additionally,  $8 \times 10^5$  pfus from the BR11 cDNA library were screened using the pooled serum described. Of these, 23 were positive.

The positive clones were subcloned, purified, and excised to forms suitable for insertion in plasmids. Following amplification of the plasmids, DNA inserts were evaluated via restriction mapping (EcoRI-XbaI), and clones which represented different cDNA inserts were sequenced using standard methodologies.

If sequences were identical to sequences found in GenBank, they were classified as known genes, while sequences which shared identity only with ESTs, or were identical to nothing in these data bases, were designated as unknown genes. Of the clones from the breast cancer library which were positive with autologous serum, 3 were unknown genes. Of the

remaining 35, 15 were identical to either NY-ESO-1, or SSX2, two known members of the CT antigen family described supra, while the remaining clones corresponded to 14 known genes. Of the testicular library, 12 of the clones were SSX2.

The NY-ESO-1 antigen was not found, probably because the commercial library that was used had been size fractionated to have an average length of 1.5 kilobases, which is larger than full length NY-ESO-1 cDNA which is about 750 base pairs long.

With respect to the screening carried out with pooled, allogeneic sera, four of the clones were NY-ESO-1. No other CT antigens were identified. With the exception of NY-ESO-1, all of the genes identified were expressed universally in normal tissue.

A full listing of the isolated genes, and their frequency of occurrence follows, in tables 1, 2 and 3. Two genes were found in both the BR11 and testicular libraries, i.e., poly (ADP-ribose) polymerase, and tumor suppression gene ING1. The poly (ADP-ribose) polymerase gene has also been found in colon cancer libraries screened via SEREX, as is disclosed by Scanlan, et al., Int. J. Cancer 76: 652-58 (1998) when the genes identified in the screening of the BR11 cDNA library by autologous and allogeneic sera were compared, NY-ESO-1 and human keratin.

**Table 1. SEREX-defined genes identified by autologous screening of BR11 cDNA library**

Gene group	No. of clones	Comments	Expression
CT genes	10	NY-ESO-1	tumor, testis
	5	SSX2	tumor, testis
Non-CT genes	5	Nuclear Receptor Co-Repressor	ubiquitous
	4	Poly(ADP-ribose) polymerase	ubiquitous
	2	Adenylosuccinatelyase	ubiquitous
	2	cosmid 313 (human)	ESTs: muscle, brain, breast
	1	CD 151 (transmembrane protein)	ubiquitous

1	Human HRY Gen	RT-PCR: multiple normal tissues
1	Alanyl-t-RNA-Synthetase	ubiquitous
1	NAD(+) ADP-Ribosyltransferase	ubiquitous
1	Human keratin 10	ESTs: multiple normal tissues
1	Human EGFR kinase substrate	ubiquitous
1	<i>ING 1</i> Tumor suppressor gene	RT-PCR: multiple normal tissues
1	Unknown gene, NCI_CGAP_Pr12 cDNA clone	ESTs: pancreas, liver, spleen, uterus
1	Unknown gene	ESTs: multiple normal tissues
1	Unknown gene	RT-PCR: multiple normal tissues

**Table 2. SEREX-defined genes identified by allogeneic screening of BR11 cDNA library**

Gene group	No. of clones	Comments	Expression
CT genes	4	NY-ESO-1	tumor, testis
Non-CT genes	6	zinc-finger helicase	ESTs: brain, fetal heart, total fetus
	4	Acetoacetyl-CoA-thiolase	ubiquitous
	3	KIAA0330 gene	ESTs: multiple normal tissues
	2	U1snRNP	ubiquitous
	1	Human aldolase A	ubiquitous
	1	Retinoblastoma binding protein 6	ESTs: tonsils, fetal brain, endothelial cells, brain
	1	$\alpha$ 2-Macroglobulin receptor associated protein	ubiquitous
	1	Human Keratin 10	ESTs: multiple normal tissues

**Table 3. SEREX-defined genes identified by screening of a testicular cDNA library with BR11 serum**

Gene group	No. of clones	Comments	Expression
CT genes:	12	SSX2	tumor, testis
Non-CT genes:	3	Rho-associated coiled-coil forming protein	ubiquitous
	3	Poly(ADP-ribose) polymerase	ubiquitous



3	Gene from HeLa cell, similar to TITIN	ubiquitous
2	Gene from parathyroid tumor	RT-PCR: multiple normal tissues
1	Transcription termination factor I-interacting peptide 21	ubiquitous
1	Gene from fetal heart	ESTs: multiple normal tissues
1	<i>ING 1</i> tumor suppressor gene	RT-PCR: multiple normal tissues
1	KIAA0647 cDNA	ESTs: multiple normal tissues
1	KIAA0667 cDNA	ESTs: multiple normal tissues

## EXAMPLE 2

The mRNA expression pattern of the cDNAs identified in example 1, in both normal and malignant tissues, was studied. To do this, gene specific oligonucleotide primers were designed which would amplify cDNA segments 300-600 base pairs in length, using a primer melting temperature of 65-70° C. The primers used for amplifying MAGE-1,2,3 and 4, BAGE, NY-ESO-1, SCP1, and SSX1, 2, 3, 4 and 5 were known primers, or were based on published sequences. See Chen, et al. *supra*; Tureci, et al., Proc. Natl. Acad. Sci. USA 95: 5211-16 (1998). Gure, et al., Int. J. Cancer 72: 965-71 (1997); Chen, et al., Proc. Natl. Acad. Sci. USA 91: 1004-1008 (1994); Gaugler, et al., J. Exp. Med. 179: 921-930 (1994), dePlaen, et al., Immunogenetics 40: 360-369 (1994), all of which are incorporated by reference. RT-PCR was carried out for 35 amplification cycles, at an annealing temperature of 60° C. Using this RT-PCR assay, the breast cancer tumor specimen was positive for a broad range of CT antigens, including MAGE-1,3 AND 4, BAGE, SSX2, NY-ESO-1 and CT7. The known CT antigens SCP-1, SSX1, 4 and 5 were not found to be expressed.

An additional set of experiments were carried out, in which the seroreactivity of patient sera against tumor antigens was tested. Specially, ELISAs were carried out, in accordance with

Stockert, et al., J. Exp. Med. 187: 1349-1354 (1998), incorporated by reference, to determine if antibodies were present in the patient sera. Assays were run for MAGE-1, MAGE-3, NY-ESO-1, and SSX2. The ELISAs were positive for NY-ESO-1 and SSX2, but not the two MAGE antigens.

### EXAMPLE 3

Two clones (one from the breast cancer cDNA library and one from the testicular library), were identified as a gene referred to as ING1, which is a tumor suppressor gene candidate. See Garkavtsev, et al., Nature 391: 295-8 (1998), incorporated by reference. The sequence found in the breast cancer library, differed from the known sequence of ING1 at six residues, i.e., positions 818, 836, 855, 861, 866 and 874. The sequence with the six variants is set forth at SEQ ID NO: 1. The sequence of wild type ING1 is set out at SEQ ID NO: 2.

To determine if any of these differences represented a mutation in tumors, a short, PCR fragment which contained the six positions referred to supra was amplified from a panel of allogeneic normal tissue, subcloned, amplified, and sequenced following standard methods.

The results indicated that the sequences in the allogeneic tissues were identical to what was found in tumors, ruling out the hypothesis that the sequence differences were a tumor associated mutation. This conclusion was confirmed, using the testicular library clone, and using restriction analysis of ING1 cDNA taken from normal tissues. One must conclude, therefore, that the sequence information provided by Garkavtsev, et al., supra, is correct.

**EXAMPLE 4**

Additional experiments were carried out to determine whether genetic variations might exist in the 5' portion of the ING1 gene, which might differ from the 5' portion of the clone discussed supra (SEQ ID NO: 1). In a first group of experiments, attempts were made to obtain full length ING1 cDNA from both the breast tumor library, and the testicular library. SEQ ID NO: 1 was used as a probe of the library, using standard methods.

Four clones were isolated from the testicular library and none were isolated from the breast cancer library. The four clones, following sequencing, were found to derive from three transcript variants. The three variants were identical from position 586 down to their 3' end, but differed in their 5' regions, suggesting alternatively spliced variants, involving the same exon-intron junction. All three differed from the sequence of ING1 described by Garkavtsev, et al., in Nat. Genet.14: 415-420 (1996). These three variants are set out as SEQ ID NOS: 1, 3 and 4.

All of the sequences were then analyzed. The ORFs of SEQ ID NOS: 2, 1 and 4 (SEQ ID NO: 2 is the originally disclosed, ING1 sequence), encode polypeptides of 294, 279 and 235 amino acids, of which 233 are encoded by the 3' region common to the three sequences. These putative sequences are set out as SEQ ID NOS:19, 5, and 7. With respect to SEQ ID NO: 3, however, no translational initiation site could be identified in its 5' region.

**EXAMPLE 5**

The data regarding SEQ ID NO: 3, described supra, suggested further experiments to find additional ORFs in the 5-end of variant transcripts of the molecule. In order to determine this, 5'-RACE -PCR was carried out using gene specific and adapted specific primers, together with commercially available products, and standard methodologies.

The primers used for these experiments were:

CACACAGGATCCATGTTGAGTCCTGCCAACGG

CGTGGTCGTGGTTGCTGGACGCG

(SEQ ID NOS: 9 and 10), for SEQ ID NO: 1;

CCCAGCGGCCCTGACGCTGTC

CGTGGTCGTGGTTGCTGGACGCG

(SEQ ID NOS: 11 and 12), for SEQ ID NO: 3; and

GGAAGAGATAAGGCCTAGGGAAG

CGTGGTCGTGGTTGCTGGACGCG

(SEQ ID NOS: 13 and 14), for SEQ ID NO: 4.

Cloning and sequencing of the products of RACE PCR showed that the variant sequence of SEQ ID NO: 4 was 5' to SEQ ID NO: 3, and that full length cDNA for the variant SEQ ID NO: 3 contained an additional exon 609 nucleotides long, positioned between SEQ ID NO: 3 and the shared, 3' sequence referred to supra. This exon did not include an ORF. The first available initiation site would be an initial methionine at amino acid 70 of SEQ ID NO: 1. Thus, if expressed, SEQ ID NO: 3 would correspond to a molecule with a 681 base pair, untranslated 5' end and a region encoding 210 amino acids (SEQ ID NO:6).

#### EXAMPLE 6

The presence of transcript variants with at least 3 different transcriptional initiation sites, and possibly different promoters, suggested that mRNA expression might be under different, tissue specific regulation.

To determine this, variant-specific primers were synthesized, and RT-PCR was carried out on a panel of tissues, using standard methods.

SEQ ID NO: 1 was found to be expressed universally in all of the normal breast, brain and testis tissues examined, in six breast cancer lines, and 8 melanoma cell lines, and in cultured melanocytes. SEQ ID NO: 3 was found to be expressed in four of the six breast cancer lines, normal testis, liver, kidney, colon and brain. SEQ ID NO: 4 was only found to be expressed by normal testis cells and weakly in brain cells.

#### EXAMPLE 7

A further set of experiments were carried out to determine if antibodies against ING1 were present in sera of normal and cancer patients. A phase plaque immuno assay of the type described supra was carried out, using clones of SEQ ID NO: 1 as target. Of 14 allogeneic sera taken from breast cancer patients, two were positive at 1:200 dilutions. All normal sera were negative.

#### EXAMPLE 8

The BR11 cDNA library described supra was then screened, using SEQ ID NO: 1 and standard methodologies. A 593 base pair cDNA was identified, which was different from any sequences in the data banks consulted. The sequence of this cDNA molecule is set out at SEQ ID NO: 8.

The cDNA molecule set forth as SEQ ID NO:1 was then used in Southern blotting experiments. In brief, genomic DNA was isolated from normal human tissue, digested with BamHI or Hind III, and then separated onto 0.7% agarose gel, blotted onto nitrocellulose filters,

and hybridized using  $^{32}\text{P}$  labelled SEQ ID NO: 1, at high stringency conditions (aqueous buffer,  $65^\circ\text{C}$ ). The probes were permitted to hybridize overnight, and then exposed for autoradiography. Two hybridizing DNA species were identified, i.e., SEQ ID NOS: 1 and 8.

#### EXAMPLE 9

The cDNA molecule set forth in SEQ ID NO: 8 was then analyzed. 5'-RACE PCR was carried out using normal fetus cDNA. Full length cDNA for the molecule is 771 base pairs long, without the poly A tail. It shows strong homology to SEQ ID NO: 1, with the strongest homology in the 5' two-thirds (76% identity over nucleotide 1-480); however, the longest ORF is only 129 base pairs, and would encode a poly peptide 42 amino acids long which was homologous to, but much shorter than, the expected expression product of SEQ ID NO: 1.

In addition to the coding region, SEQ ID NO: 8 contains 203 base pairs of 5'-untranslated region, and 439 base pairs of 3'-untranslated region.

RT-PCR assays were carried out, as described supra. All of the normal tissues tested, including brain, colon, testis, tissue and breast, were positive for expression of this gene. Eight melanoma cell lines were tested, of which seven showed varying levels of expression, and one showed no expression. Six breast cancer cell lines were tested, of which four showed various levels of expression, and two showed no expression.

#### EXAMPLE 10

An additional breast cancer cDNA library, referred to as "BR17-128", was screened, using autologous sera. A cDNA molecule was identified.

Analysis of the sequence suggested that it was incomplete at the 5' end. To extend the sequence, a testicular cDNA library was screened with a nucleotide probe based upon the partial sequence identified in the breast cancer library. An additional 1200 base pairs were identified following these screenings. The 2011 base pairs of information are set forth in SEQ ID NO: 15.

The longest open reading frame is 1539 base pairs, corresponding to a protein of about 59.15 kilodaltons. The deduced sequence is set forth at SEQ ID NO: 16.

RT-PCR was then carried out using the following primers:

CACACAGGATCCATGCAGGCCCGCACAAGGAG

CACACAAAGCTTCTAGGATTTGGCACAGCCAGAG

(SEQ ID NOS: 17 and 18)

Strong signals were observed in normal testis and breast tissue, and weak expression was observed in placenta.

No expression was found in normal brain, kidney, liver, colon, adrenal, fetal brain, lung, pancreas, prostate, thymus, uterus, and ovary tissue of tumor cell lines tested, 2 of the breast cancer lines were strongly positive and two were weakly positive. Of melanoma two of 8 were strongly positive, and 3 were weakly positive. Of lung cancer cell lines, 4 of 15 were strongly positive, and 3 were weakly positive.

When cancer tissue specimens were tested, 16 of 25 breast cancer samples were strongly positive, and 3 additional samples were weakly positive. Two of 36 melanoma samples were positive (one strong, one weak). All other cancer tissue samples were negative.

When Northern blotting was carried out, a high molecular weight smear was observed in testis, but in no other tissues tested.

**EXAMPLE 11**

Further experiments were carried out using the tumor sample referred to in example 10, supra. This sample was derived from a subcutaneous metastasis of a 60 year old female breast cancer patient. Total RNA was extracted, as described supra. Following the extraction, a cDNA library was constructed in  $\lambda$ -ZAP expression vectors, also as described supra. Screening was carried out, using the protocol set forth in example 1. A total of  $7 \times 10^5$  pfus were screened. Fourteen reactive clones were identified, purified, and sequenced. The sequences were then compared to published sequences in GenBank and EST databases. These analyses indicated that the clones were derived from seven distinct genes, two of which were known, and five unknown. The two known genes were "PBK-1" (three clones), and TI-227 (one clone). These are universally expressed genes, with the libraries referred to supra showing ESTs for these genes from many different tissues.

With respect to the remaining 10 clones, six were derived from the same gene, referred to hereafter as "NY-BR-1." Three cDNA sequences were found in the EST database which shared identity with the gene. Two of these (AI 951118 and AW 373574) were identified as being derived from a breast cancer library, while the third (AW 170035), was from a pooled tissue source.

**EXAMPLE 12**

The distribution of the new gene NY-BR-1 referred to supra was determined via RT-PCR. In brief, gene specific oligonucleotide NY-BR-1 primers were designed to amplify cDNA



segments 300-600 base pairs in length, with primer melting temperatures estimated at 65-70°C.

The RT-PCR was then carried out over 30 amplification cycles, using a thermal cycler, and an annealing temperature of 60°C. Products were analyzed via 1.5% gel electrophoresis, and ethidium bromide visualization. Fifteen normal tissues (adrenal gland, fetal brain, lung, mammary gland, pancreas, placenta, prostate, thymus, uterus, ovary, brain, kidney, liver, colon and testis) were assayed. The NY-BR-1 clone gave a strong signal in mammary gland and testis tissue, and a very faint signal in placenta. All other tissues were negative. The other clones were expressed universally, based upon comparison to information in the EST database library, and were not pursued further.

The expression pattern of NY-BR-1 in cancer samples was then tested, by carrying out RT-PCR, as described supra, on tumor samples.

In order to determine the expression pattern, primers:

caaagcagag                      cctcccgaga                      ag

(SEQ ID NO: 20) and

cctatgctgc                      tcttcgattc                      ttcc

(SEQ ID NO: 21) were used.

Of twenty-five breast cancer samples tested, twenty two were positive for NY-BR-1. Of these, seventeen gave strong signals, and five gave weak to modest signals.

An additional 82 non-mammary tumor samples were also analyzed, divided into 36 melanoma, 26 non small cell lung cancer, 6 colon cancer, 6 squamous cell carcinoma, 6 transitional cell carcinoma, and two leiomyosarcomas. Only two melanoma samples were positive for NY-BR-1 expression.

The study was then extended to expression of NY-BR-1 in tissue culture. Cell lines derived from breast tumor, melanoma, and small cell lung cancer were studied. Four of six breast cancer cells were positive (two were very weak), four of eight melanoma (two very weak), and seven of fourteen small cell lung cancer lines (two very weak) were positive.

### EXAMPLE 13

In order to determine the complete cDNA molecule for NY-BR-1, the sequences of the six clones referred to supra were compiled, to produce a nucleotide sequence 1464 base pairs long. Analysis of the open reading frame showed a continuous ORF throughout, indicating that the compiled sequence is not complete.

Comparison of the compiled sequence with the three EST library sequences referred to supra allowed for extension of the sequence. The EST entry AW170035 (446 base pairs long) overlapped the compiled sequence by 89 base pairs at its 5' end, permitting extension of the sequence by another 357 base pairs. A translational terminal codon was identified in this way, leading to a molecule with a 3'-untranslated region 333 base pairs long. The 5' end of the molecule was lacking, however, which led to the experiment described infra.

### EXAMPLE 14

In order to determine the missing, 5' end of the clone described supra, a commercially available testis cDNA expression library was screened, using a PCR expression product of the type described supra as a probe. In brief,  $5 \times 10^4$  pfus per 150 mm plate were transferred to nitrocellulose membranes, which were then submerged in denaturation solution (1.5M NaCl and 0.5 M NaOH), transferred to neutralization solution (1.5 M NaCl and 0.5M Tris-HCl), and then

rinsed with 0.2M Tris-HCl, and 2xSSC. Probes were labelled with  $^{32}\text{P}$  and hybridization was carried out at high stringency conditions (i.e., 68°C, aqueous buffer). Any positive clones were subcloned, purified, and in vivo excised to plasmid PBK-CMV, as described supra.

One of the clones identified in this way included an additional 1346 base pairs at the 5' end; however, it was not a full length molecule. A 5'-RACE-PCR was carried out, using commercially available products. The PCR product was cloned into plasmid vector pGEMT and sequenced. The results indicated that cDNA sequence was extended 1292 base pairs further, but no translation initiation site could be determined, because no stop codons could be detected. It could be concluded, however, that the cDNA of the NY-BR17 clone comprises at least 4026 nucleotides, which are presented as SEQ ID NO: 22. The molecule, as depicted, encodes a protein at least about 152.8 kDA in molecular weight. Structurally, there are 99 base pairs 5' to the presumed translation initiation site, and an untranslated segment 333 base pairs long at the 3' end. The predicted amino acid sequence of the coding region for SEQ ID NO: 22 is set out at SEQ. ID NO: 23.

SEQ ID NO: 23 was analyzed for motifs, using the known search programs PROSITE and Pfam. A bipartite nuclear localization signal motif was identified at amino acids 17-34, suggesting that the protein is a nuclear protein. Five tandem ankyrin repeats were identified, at amino acids 49-81, 82-114, 115-147, 148-180 and 181-213. A bZIP site (i.e. a DNA binding site followed by a leucine zipper motif) was found at amino acid positions 1077-1104, suggesting a transcription factor function. It was also observed that three repetitive elements were identified in between the ankyrin repeats and the bZIP DNA binding site. To elaborate, a repetitive element 117 nucleotides long is tandemly repeated 3 times, between amino acids 459-815. The second repetitive sequence, consisting of 11 amino acids, repeats 7 times between amino acids

224 and 300. The third repetitive element, 34 amino acids long, is repeated twice, between amino acids 301-368.

#### EXAMPLE 15

The six clones described supra were compared, and analysis revealed that they were derived from two different splice variants. Specifically, two clones, referred to as "BR17-8" and "BR 17-44a", contain one more exon, of 111 base pairs (nucleotides 3015-3125 of SEQ ID NO: 22), which encodes amino acids 973-1009 of SEQ ID NO: 23, than do clones BR 17-1a, BR17-35b and BR17-44b. The shortest of the six clones, BR17-128, starts 3' to the additional exons. The key structural elements referred to supra were present in both splice variants, suggesting that there was no difference in biological function.

The expression pattern of the two splice variants was assessed via PT-PCR, using primers which spanned the 111 base pair exon referred to supra.

The primers used were:

aatgggaaca                      agagctctgc                      ag

(SEQ ID NO: 24) and

gggtcatctg                      aagttcagca                      ttc

(SEQ ID NO: 25)

Both variants were expressed strongly in normal testis and breast. The longer variant was dominant in testis, and the shorter variant in breast cells. When breast cancer cells were tested, co-typing of the variant was observed, (7 strongly, 2 weakly positive, and 1 negative), with the shorter variant being the predominant form consistently.

**EXAMPLE 16**

The frequency of antibody response against NY-BR-1 in breast cancer patients was tested. To do this, a recombinant protein consisting of amino acids 993-1188 of SEQ ID NO: 23 was prepared. (This is the protein encoded by clone BR 17-128, referred to supra). A total of 140 serum samples were taken from breast cancer patients, as were 60 normal serum samples. These were analyzed via Western blotting, using standard methods.

Four of the cancer sera samples were positive, including a sample from patient BR17. All normal sera were negative.

An additional set of experiments was then carried out to determine if sera recognized the portion of NY-BR-1 protein with repetitive elements. To do this, a different recombinant protein, consisting of amino acids 405-1000 was made, and tested in Western blot assays. None of the four antibody positive sera reacted with this protein indicating that an antibody epitope is located in the non-repetitive, carboxy terminal end of the molecule.

**EXAMPLE 17**

The screening of the testicular cDNA library referred to supra resulted, inter alia, in the identification of a cDNA molecule that was homologous to NY-BR-1. The molecule is 3673 base pairs in length, excluding the poly A tail. This corresponded to nucleotides 1-3481 of SEQ ID NO: 22, and showed 62% homology thereto. No sequence identity to sequences in libraries was noted. ORF analysis identified an ORF from nucleotide 641 through the end of the sequence, with 54% homology to the protein sequence of SEQ. ID NO: 23. The ATG initiation codon of this sequence is 292 base pairs further 3' to the presumed initiation codon of NY-BR-1, and is preceded by 640 untranslated base pairs at its 5' end. This 640 base pair sequence includes

scattered stop codons. The nucleotide sequence and deduced amino acid sequence are presented as SEQ ID NOS: 26 and 27, respectively.

RT-PCR analysis was carried out in the same way as is described supra, using primers:

tct catagat      gctggtgctg      atc

(SEQ ID NO: 28) and

cccagacatt      gaattttggc      agac

(SEQ ID NO: 29).

Tissue restricted mRNA expression was found. The expression pattern differed from that of SEQ ID NO: 22. In brief, of six normal tissues examined, strong signals were found in brain and testis only. There was no or weak expression in normal breast tissues, and kidney, liver and colon tissues were negative. Eight of ten 10 breast cancer specimens tested supra were positive for SEQ. ID NO: 26. Six samples were positive for both SEQ. ID NO: 22 and 26, one for SEQ. ID NO: 22 only, two for the SEQ. ID NO: 26 only, and one was negative for both.

## EXAMPLE 18

Recently, a working draft of the human genome sequence was released. This database was searched, using standard methods, and NY-BR-1 was found to have sequence identity with at least three chromosome 10 clones, identified by Genbank accession numbers AL157387, AL37148, and AC067744. These localize NY-BR-1 to chromosome 10 p11.21-12.1.

The comparison of NY-BR-1 and the human genomic sequence led to definition of NY-BR-1 exon-intron organization. In brief, the coding region of the gene contains essentially 19 structurally distinct exons with at least 2 exons encoding 3' untranslated regions. Detailed exon-intron junction information is described at Genbank AF 269081.

The six ankyrin repeats, referred to supra, are all found within exon 7. The 357 nucleotide repeating unit is composed of exons 10-15. The available genomic sequences are not complete, however, and only one of the three copies was identified, suggesting that DNA sequences between exons 5 and 10 may be duplicated and inserted in tandem, during genetic evolution. In brief, when the isolated NY-BR-1 cDNA clone was analyzed, three complete and one incomplete copy of the repeating units are present. The exon sequences can be expressed as exons 1-2-3-4-5-6-7-8-9-(10-11-12-13-14-15)-(10A-11A-12A-13A-14A-15A)-(10B-11B-12B-13B-14B-15B)-(10C-11C-12C-13C-14C)-16-17-18-19-20-21, wherein A, B & C are inexact copies of exon 10-15 sequences. Cloned, NY-BR-1 cDNA has 38 exons in toto.

It was noted, supra, that the sequence of NY-BR-1 cDNA was not complete at the 5' end. Genomic sequence (Genbank AC067744), permitted extension of the 5' end. Translation of the 5' genomic sequence led to the identification of a new translation initiation site, 168 base pairs upstream of the previously predicted ATG initiation codon. This led to an NY-BR-1 polypeptide including 1397 amino acid longer, 56 residue of which are added at the N-terminus, compared to prior sequence information, i.e.:

MEEISAAAVKVVPGERPSPFSQLVYTSNDSYIVHSGDLRKIHKAASRGQVRKLEK (SEQ ID NO: 30).

## EXAMPLE 20

Reference was made, supra, to the two difference splice variants of NY-BR-1. Comparison of the splice variants with the genomic sequence confirmed that an alternate splicing event, with the longer variant incorporating part of intron 33 into exon 34 (i.e., exon 17 of the basic exon/intron framework described supra).

Key structural elements that were predicted in NY-BR-1, described supra, are present in both variants, suggesting that there is no difference in biological function, or subcellular location.

#### EXAMPLE 21

As with NY BR-1, the variant NY-BR-1.1, described supra, was screened against the working draft of the human genome sequence. One clone was found with sequence identity, i.e., GenBank AL359312, derive from chromosome 9. Thus, NY-BR-1 and NY-BR-1.1 both appear to be functioning genes, on two different chromosomes. The Genbank sequence referred to herein does not contain all of NY-BR-1.1, which precludes defining exon-intron structure. Nonetheless, at least 3 exons can be defined, which correspond to exons 16-18 of the NY-BR-1 basic framework. Exon-intron junctions are conserved.

#### EXAMPLE 22

A series of peptides were synthesized, based upon the amino acid sequence of NY-BR-1, as set forth in SEQ ID NO: 23. These were then tested for their ability to bind to HLA-A2 molecules and to stimulate CTL proliferation, using an ELISPOT assay. This assay involved coating 96-well, flat bottom nitrocellulose plates with 5ug/ml of anti-interferon gamma antibodies in 100 ul of PBS per well, followed by overnight incubation. Purified CD8<sup>+</sup> cells, which had been separated from PBL samples via magnetic beads coated with anti-CD8 antibodies were then added, at  $1 \times 10^5$  cells/well, in RPMI 1640 medium, that had been supplemented with 10% human serum, L-asparagine (50 mg/l), L-arginine (242 mg/l), L-glutamine (300 mg/l), together with IL-2 (2.5ng/ml), in a final volume of 100 ul. CD8<sup>+</sup> effector



cells were prepared by presensitizing with peptide, and were then added at from  $5 \times 10^3$  to  $2 \times 10^4$  cells/well. Peptides were pulsed onto irradiated T2 cells at a concentration of 10ug/ml for 1 hour, washed and added to effector cells, at  $5 \times 10^4$  cells/well. The plates were incubated for 16 hours at 37°C, washed six times with 0.05% Tween 20/PBS, and were then supplemented with biotinylated, anti-interferon gamma specific antibody at 0.5 ug/ml. After incubation for 2 hours at 37°C, plates were washed, and developed with commercially available reagents, for 1 hour, followed by 10 minutes of incubation with dye substrate. Plates were then prepped for counting, positives being indicated by blue spots. The number of blue spots/well was determined as the frequency of NY-ESO-1 specific CTLs/well.

Experiments were run, in triplicate, and total number of CTLs was calculated. As controls, one of reagents alone, effector cells alone, or antigen presenting cells alone were used. The difference between the number of positives in stimulated versus non-stimulated cells, was calculated as the effective number of peptide specific CTLs above background. Three peptides were found to be reactive, i.e.:

LLSHGAVIEV (amino acids 102-111 of SEQ ID NO: 23)

SLSKILDTV (amino acids 904-912 of SEQ ID NO: 23 )

SLDQKLFQL (amino acids 1262-1270 of SEQ ID NO: 23).

The complete list of peptides tested, with reference to their position in SEQ ID NO: 23, follows:

Peptide	Position
FLVDRKVCQL	35-43
ILIDSGADI	68-76

AVYSEILSV	90-98
ILSVVAKLL	95-103
LLSHGAVIEV	102-111
KLLSHGAVI	101-109
FLLIKNANA	134-142
MLLQQNVDV	167-175
GMLLQQNVDV	166-175
LLQQNVDVFA	168-177
IAWEKKETPV	361-370
SLFESSAKI	430-438
CIPENSIYQKV	441-450
KVMEINREV	449-457
ELMDMQTFKA	687-696
ELMDMQTFKA	806-815
SLSKILDTV	904-912
KILDTVHSC	907-915
ILNEKIREEL	987-996
RIQDIELKSV	1018-1027
YLLHENCML	1043-1051
CMLKKEIAML	1049-1058
AMLKLELATL	1056-1065
KILKEKNAEL	1081-1090
VLIAENTML	1114-1122
CLQRKMNV DV	1174-1183
KMNV DV SST	1178-1186
SLDQKLFQL	1262-1270
KLFQLQSKNM	1266-1275

FQLQSKNMWL	1268-1277
QLQSKNMWL	1269-1277
NMWLQQQLV	1274-1282
WLQQQLVHA	1276-1284
KITIDIHFL	1293-1301

The foregoing examples describe the isolation of a nucleic acid molecule which encodes a cancer associated antigen. "Associated" is used herein because while it is clear that the relevant molecule was expressed by several types of cancer, other cancers, not screened herein, may also express the antigen.

The invention relates to nucleic acid molecules which encode the antigens encoded by, e.g., SEQ ID NOS: 1, 3, 8, 15, 22 and 26 as well as the antigens encoded thereby, such as the proteins with the amino acid sequences of SEQ ID NOS: 5, 6, 7, 16, 23, 27, and 30. It is to be understood that all sequences which encode the recited antigen are a part of the invention.

Also a part of the invention are proteins, polypeptides, and peptides, which comprise, e.g., at least nine consecutive amino acids found in SEQ ID NO: 23, or at least nine consecutive amino acids of the amino acids of SEQ ID NO: 30. Proteins, polypeptides and peptides comprising nine or more amino acids of SEQ ID NO: 5, 6, 7, 16 or 27 are also a part of the invention. Especially preferred are peptides comprising or consisting of amino acids 102-111, 904-912, or 1262-1270 of SEQ ID NO: 23. Such peptides may, but do not necessarily provoke CTL responses when complexed with an HLA molecule, such as an HLA-A2 molecule. They may also bind to different MHC or HLA molecules, including, but not being limited to, HLA-A1, A2, A3, B7, B8, Cw3, Cw6, or serve, e.g., as immunogens, as part of immunogenic cocktail compositions, where they are combined with other proteins or polypeptides, and so forth. Also

a part of the invention are the nucleic acid molecules which encode these molecules, such as "minigenes," expression vectors that include the coding regions, recombinant cells containing these, and so forth. All are a part of the invention.

Also a part of the invention are expression vectors which incorporate the nucleic acid molecules of the invention, in operable linkage (i.e., "operably linked") to a promoter. Construction of such vectors, such as viral (e.g., adenovirus or Vaccinia virus) or attenuated viral vectors is well within the skill of the art, as is the transformation or transfection of cells, to produce eukaryotic cell lines, or prokaryotic cell strains which encode the molecule of interest. Exemplary of the host cells which can be employed in this fashion are COS cells, CHO cells, yeast cells, insect cells (e.g., Spodoptera frugiperda), NIH 3T3 cells, and so forth. Prokaryotic cells, such as E. coli and other bacteria may also be used. Any of these cells can also be transformed or transfected with further nucleic acid molecules, such as those encoding cytokines, e.g., interleukins such as IL-2, 4, 6, or 12 or HLA or MHC molecules.

Also a part of the invention are the antigens described herein, both in original form and in any different post translational modified forms. The molecules are large enough to be antigenic without any posttranslational modification, and hence are useful as immunogens, when combined with an adjuvant (or without it), in both precursor and post-translationally modified forms. Antibodies produced using these antigens, both poly and monoclonal, are also a part of the invention as well as hybridomas which make monoclonal antibodies to the antigens. The whole protein can be used therapeutically, or in portions, as discussed infra. Also a part of the invention are antibodies against this antigen, be these polyclonal, monoclonal, reactive fragments, such as Fab, (F(ab)<sub>2</sub>)' and other fragments, as well as chimeras, humanized antibodies, recombinantly produced antibodies, and so forth.


As is clear from the disclosure, one may use the proteins and nucleic acid molecules of the invention diagnostically. The SEREX methodology discussed herein is premised on an immune response to a pathology associated antigen. Hence, one may assay for the relevant pathology via, e.g., testing a body fluid sample of a subject, such as serum, for reactivity with the antigen per se. Reactivity would be deemed indicative of possible presence of the pathology. So, too, could one assay for the expression of any of the antigens via any of the standard nucleic acid hybridization assays which are well known to the art, and need not be elaborated upon herein. One could assay for antibodies against the subject molecules, using standard immunoassays as well.

Analysis of SEQ ID NO: 1, 3, 4, 8, 15, 22 and 26 will show that there are 5' and 3' non-coding regions presented therein. The invention relates to those isolated nucleic acid molecules which contain at least the coding segment, and which may contain any or all of the non-coding 5' and 3' portions.

Also a part of the invention are portions of the relevant nucleic acid molecules which can be used, for example, as oligonucleotide primers and/or probes, such as one or more of SEQ ID NOS: 9, 10, 11, 12, 13, 14, 17, 18, 20, 21, 24, 25, 28, and 29 as well as amplification products like nucleic acid molecules comprising at least nucleotides 305-748 of SEQ ID NO: 1, or amplification products described in the examples, including those in examples 12, 14, etc.

As was discussed supra, study of other members of the "CT" family reveals that these are also processed to peptides which provoke lysis by cytolytic T cells. There has been a great deal of work on motifs for various MHC or HLA molecules, which is applicable here. Hence, a further aspect of the invention is a therapeutic method, wherein one or more peptides derived from the antigens of the invention which bind to an HLA molecule on the surface of a patient's

tumor cells are administered to the patient, in an amount sufficient for the peptides to bind to the MHC/HLA molecules, and provoke lysis by T cells. Any combination of peptides may be used. These peptides, which may be used alone or in combination, as well as the entire protein or immunoreactive portions thereof, may be administered to a subject in need thereof, using any of the standard types of administration, such as intravenous, intradermal, subcutaneous, oral, rectal, and transdermal administration. Standard pharmaceutical carriers, adjuvants, such as saponins, GM-CSF, and interleukins and so forth may also be used. Further, these peptides and proteins may be formulated into vaccines with the listed material, as may dendritic cells, or other cells which present relevant MHC/peptide complexes.

Similarly, the invention contemplates therapies wherein nucleic acid molecules which encode the proteins of the invention, one or more or peptides which are derived from these proteins are incorporated into a vector, such as a Vaccinia or adenovirus based vector, to render it transfectable into eukaryotic cells, such as human cells. Similarly, nucleic acid molecules which encode one or more of the peptides may be incorporated into these vectors, which are then the major constituent of nucleic acid bases therapies. 

Any of these assays can also be used in progression/regression studies. One can monitor the course of abnormality involving expression of these antigens simply by monitoring levels of the protein, its expression, antibodies against it and so forth using any or all of the methods set forth supra.

It should be clear that these methodologies may also be used to track the efficacy of a therapeutic regime. Essentially, one can take a baseline value for a protein of interest using any of the assays discussed supra, administer a given therapeutic agent, and then monitor levels of the protein thereafter, observing changes in antigen levels as indicia of the efficacy of the regime.

As was indicated supra, the invention involves, inter alia, the recognition of an “integrated” immune response to the molecules of the invention. One ramification of this is the ability to monitor the course of cancer therapy. In this method, which is a part of the invention, a subject in need of the therapy receives a vaccination of a type described herein. Such a vaccination results, e.g., in a T cell response against cells presenting HLA/peptide complexes on their cells. The response also includes an antibody response, possibly a result of the release of antibody provoking proteins via the lysis of cells by the T cells. Hence, one can monitor the effect of a vaccine, by monitoring an antibody response. As is indicated, supra, an increase in antibody titer may be taken as an indicia of progress with a vaccine, and vice versa. Hence, a further aspect of the invention is a method for monitoring efficacy of a vaccine, following administration thereof, by determining levels of antibodies in the subject which are specific for the vaccine itself, or a large molecule of which the vaccine is a part.

The identification of the subject proteins as being implicated in pathological conditions such as cancer also suggests a number of therapeutic approaches in addition to those discussed supra. The experiments set forth supra establish that antibodies are produced in response to expression of the protein. Hence, a further embodiment of the invention is the treatment of conditions which are characterized by aberrant or abnormal levels of one or more of the proteins, via administration of antibodies, such as humanized antibodies, antibody fragments, and so forth. These may be tagged or labelled with appropriate cystostatic or cytotoxic reagents.

T cells may also be administered. It is to be noted that the T cells may be elicited in vitro using immune responsive cells such as dendritic cells, lymphocytes, or any other immune responsive cells, and then reperused into the subject being treated.

Note that the generation of T cells and/or antibodies can also be accomplished by administering cells, preferably treated to be rendered non-proliferative, which present relevant T cell or B cell epitopes for response, such as the epitopes discussed supra.

The therapeutic approaches may also include antisense therapies, wherein an antisense molecule, preferably from 10 to 100 nucleotides in length, is administered to the subject either "neat" or in a carrier, such as a liposome, to facilitate incorporation into a cell, followed by inhibition of expression of the protein. Such antisense sequences may also be incorporated into appropriate vaccines, such as in viral vectors (e.g., Vaccinia), bacterial constructs, such as variants of the known BCG vaccine, and so forth.

Other features and applications of the invention will be clear to the skilled artisan, and need not be set forth herein. The terms and expression which have been employed are used as terms of description and not of limitation, and there is no intention in the use of such terms and expression of excluding any equivalents of the features shown and described or portions thereof, it being recognized that various modifications are possible within the scope of the invention.



We claim:

1. Isolated nucleic acid molecule which encodes a cancer associated antigen, whose amino acid sequence is identical to the amino acid sequence encoded by the nucleotide sequence of SEQ ID NO: 1, 3, 4, 8, 15, 19, 22, or 26.
2. The isolated nucleic acid molecule of claim 1, comprising the nucleotide sequence of SEQ ID NO: 1.
3. The isolated nucleic acid molecule of claim 1, comprising the nucleotide sequence of SEQ ID NO: 3.
4. The isolated nucleic acid molecule of claim 1, comprising the nucleotide sequence of SEQ ID NO: 4.
5. The isolated nucleic acid molecule of claim 1, comprising the nucleotide sequence of SEQ ID NO: 8.
6. The isolated nucleic acid molecule of claim 1, comprising the nucleotide sequence of SEQ ID NO: 15.
7. The isolated nucleic acid molecule of claim 1, comprising the nucleotide sequence of SEQ ID NO: 19.
8. The isolated nucleic acid molecule of claim 1, comprising the nucleotide sequence of SEQ ID NO: 22.
9. The isolated nucleic acid molecule of claim 1, comprising the nucleotide sequence of SEQ ID NO: 26.
10. Expression vector comprising the isolated nucleic acid molecule of claim 1, operably linked to a promoter.

11. Eukaryotic cell line or prokaryotic cell strain, transformed or transfected with the expression vector of claim 10.
12. Isolated cancer associated antigen comprising all or part of the amino acid sequence encoded by SEQ ID NO: 1, 3, 4, 8, 15, 19, 22 or 26.
13. Eukaryotic cell line or prokaryotic cell strain, transformed or transfected with the isolated nucleic acid molecule of claim 1.
14. The eukaryotic cell line or prokaryotic cell strain of claim 13, wherein said cell line is also transfected with a nucleic acid molecule coding for a cytokine.
15. The eukaryotic cell line or prokaryotic cell strain of claim 14, wherein said cell line is further transfected by a nucleic acid molecule coding for an MHC molecule.
16. The eukaryotic cell line or prokaryotic cell strain of claim 14, wherein said cytokine is an interleukin.
17. The eukaryotic cell line or prokaryotic cell strain of claim 16, wherein said interleukin is IL-2, IL-4 or IL-12.
18. The eukaryotic cell line or prokaryotic cell strain of claim 13, wherein said cell line has been rendered non-proliferative.
19. The eukaryotic cell line of claim 13, wherein said cell line is a fibroblast cell line.
20. Expression vector comprising a mutated or attenuated virus and the isolated nucleic acid molecule of claim 1.
21. The expression vector of claim 20, wherein said virus is adenovirus or vaccinia virus.
22. The expression vector of claim 21, wherein said virus is vaccinia virus.
23. The expression vector of claim 21, wherein said virus is adenovirus.

24. Expression system useful in transfecting a cell, comprising (i) a first vector containing a nucleic acid molecule which codes for the isolated cancer associated antigen of claim 13 and (ii) a second vector selected from the group consisting of (a) a vector containing a nucleic acid molecule which codes for an MHC or HLA molecule which presents an antigen derived from said cancer associated antigen and (b) a vector containing a nucleic acid molecule which codes for an interleukin.

25. Immunogenic composition comprising the isolated cancer antigen of claim 12, and a pharmaceutically acceptable adjuvant.

26. The immunogenic composition of claim 25, wherein said adjuvant is a cytokine, a saponin, or GM-CSF.

27. Immunogenic composition comprising at least one peptide consisting of an amino acid sequence of from 8 to 12 amino acids concatenated to each other in the isolated cancer associated cancer antigen of claim 12, and a pharmaceutically acceptable adjuvant.

28. The immunogenic composition of claim 27, wherein said adjuvant is a saponin, a cytokine, or GM-CSF.

29. The immunogenic composition of claim 25, wherein said composition comprises a plurality of peptides which complex with a specific MHC molecule.

30. Immunogenic composition which comprises at least one expression vector which encodes a peptide derived from the amino acid sequence encoded by SEQ ID NO: 1, 3, 4, 8, 15, 19, 22 or 26.

31. The immunogenic composition of claim 30, wherein said at least one expression vector codes for a plurality of peptides.

32. Vaccine useful in treating a subject afflicted with a cancerous condition comprising the isolated eukaryotic cell line of claim 13 and a pharmacologically acceptable adjuvant.

33. The vaccine of claim 32, wherein said eukaryotic cell line has been rendered non-proliferative.

34. The vaccine of claim 33, wherein said eukaryotic cell line is a human cell line.

35. A composition of matter useful in treating a cancerous condition comprising a non-proliferative cell line having expressed on its surface a peptide derived from the amino acid sequence encoded by SEQ ID NO: 1, 3, 4, 8, 15, 19, 22 or 26.

36. The composition of matter of claim 35, wherein said cell line is a human cell line.

37. A composition of matter useful in treating a cancerous condition, comprising (i) a peptide derived from the amino acid sequence encoded by SEQ ID NO: 1, 3, 4, 8, 15, 19, 22 or 26, (ii) an MHC or HLA molecule, and (iii) a pharmaceutically acceptable carrier.

38. Isolated antibody which is specific for the cancer associated antigen of claim 12.

39. The isolated antibody of claim 38, wherein said antibody is a monoclonal antibody.

40. Method for screening for cancer in a sample, comprising contacting said sample with a nucleic acid molecule which hybridizes to all or part of the molecule encoded by SEQ ID NO: 1, 2, 3, 4, 8, 15, 19, 22 or 26 and determining hybridization as an indication of cancer cells in said sample.

41. A method for screening for cancer in a sample, comprising contacting said sample with the isolated antibody of claim 38, and determining binding of said antibody to a target as an indicator of cancer.

42. Method for diagnosing a cancerous condition in a subject, comprising contacting an immune reactive cell containing sample of said subject to a cell line transfected with the isolated nucleic acid molecule of claim 1, and determining interaction of said transfected cell line with said immunoreactive cell, said interaction being indicative of said cancer condition.

43. A method for determining regression, progression of onset of a cancerous condition comprising monitoring a sample from a patient with said cancerous condition for a parameter selected from the group consisting of (i) a protein encoded by SEQ ID NO: 1, 2, 3, 4, 8, 15, 19, 22 or 26, (ii) a peptide derived from said protein, (iii) cytolytic T cells specific for said peptide and an MHC molecule with which it non-covalently complexes, and (iv) antibodies specific for said CT protein, wherein amount of said parameter is indicative of progression or regression or onset of said cancerous condition.

44. The method of claim 43, wherein said sample is a body fluid or exudate.

45. The method of claim 43, wherein said sample is a tissue.

46. The method of claim 43, comprising contacting said sample with an antibody which specifically binds with said protein or peptide.

47. The method of claim 46, wherein said antibody is labelled with a radioactive label or an enzyme.

48. The method of claim 46, wherein said antibody is a monoclonal antibody.

49. The method of claim 43, comprising amplifying RNA which codes for said protein.

50. The method of claim 49, wherein said amplifying comprises carrying out polymerase chain reaction.

51. The method of claim 42, comprising contacting said sample with a nucleic acid molecule which specifically hybridizes to a nucleic acid molecule which codes for or expresses said protein.

52. The method of claim 49, wherein said nucleic acid molecule comprises SEQ ID NO: 9, 10, 11, 12, 13, 14, 17, 18, 20, 21, 24, 25, 28 or 29.

53. The method of claim 43, comprising assaying said sample for shed protein.

54. The method of claim 43, comprising assaying said sample for antibodies specific for said protein, by contacting said sample with protein.

55. Method for diagnosing a cancerous condition comprising assaying a sample taken from a subject for an immunoreactive cell specific for a peptide derived from a protein encoded by SEQ ID NO: 1, 2, 3, 4, 8, 15, 19, 22 or 26, complexed to an MHC molecule, presence of said immunoreactive cell being indicative of said cancerous condition.

56. Composition comprising at least one peptide consisting of an amino acid sequence of from 8 to 25 amino acids concatenated to each other in the isolated cancer associated antigen of claim 12, and a pharmaceutically acceptable adjuvant.

57. The composition of claim 56, wherein said adjuvant is a saponin, a cytokine, or GM-CSF.

58. The composition of claim 56, comprising a plurality of MHC binding peptides.

59. Composition comprising an expression vector which encodes at least one peptide consisting of an amino acid sequence of from 8 to 25 amino acids concatenated to each other in the isolated cancer associated antigen of claim 12, and pharmaceutically acceptable adjuvant.

60. The composition of claim 59, wherein said expression vector encodes a plurality of peptides.

61. A method for screening for possible presence of a pathological condition, comprising assaying a sample from a patient believed to have a pathological condition for antibodies specific to at least one of the cancer associated antigens encoded by SEQ ID NOS: 1, 2, 3, 4, 8, 15, 19, 22 or 26, presence of said antibodies being indicative of possible presence of said pathological condition.

62. The method of claim 61, wherein said pathological condition is cancer.

63. The method of claim 61, wherein said cancer is melanoma.

64. The method of claim 61, further comprising contacting said sample to purified cancer associated antigen encoded by SEQ ID NO: 1, 3, 4, 8, 15, 19, 22 or 26.

65. A method for screening for possible presence of a pathological condition in a subject, comprising assaying a sample taken from said subject for expression of a nucleic acid molecule, the nucleotide sequence of which comprises SEQ ID NO: 1, 2, 3, 4, 8, 15, 19, 22 or 26, expression of said nucleic acid molecule being indicative of possible presence of said pathological condition.

66. The method of claim 65, wherein said pathological condition is cancer.

67. The method of claim 65, comprising determining expression via polymerase chain reaction.

68. The method of claim 65, comprising determining expression by contacting said sample with at least one of SEQ ID NO: 9, 10, 11, 12, 13, 14, 17, 18, 20, 21, 24, 25, 28 or 29.

69. A method for determining regression, progression of onset of a cancerous condition comprising monitoring a sample from a patient with said cancerous condition for a parameter selected from the group consisting of (i) a cancer associated antigen encoded by SEQ ID NO: 1, 2, 3, 4, 8, 15, 19, 22 or 25, (ii) a peptide derived from said cancer associated antigen,

(iii) cytolytic T cells specific for said peptide and an MHC molecule with which it non-covalently complexes, and (iv) antibodies specific for said cancer associated antigen, wherein amount of said parameter is indicative of progression or regression or onset of said cancerous condition.

70. The method of claim 69, wherein said sample is a body fluid or exudate.

71. The method of claim 69, wherein said sample is a tissue.

72. The method of claim 69, comprising contacting said sample with an antibody which specifically binds with said protein or peptide.

73. The method of claim 72, wherein said antibody is labelled with a radioactive label or an enzyme.

74. The method of claim 72, wherein said antibody is a monoclonal antibody.

75. The method of claim 69, comprising amplifying RNA which codes for said protein.

76. The method of claim 75, wherein said amplifying comprises carrying out polymerase chain reaction.

77. The method of claim 69, comprising contacting said sample with a nucleic acid molecule which specifically hybridizes to a nucleic acid molecule which codes for or expresses said protein.

78. The method of claim 69, comprising assaying said sample for shed cancer associated antigen.

79. The method of claim 69, comprising assaying said sample for antibodies specific for said cancer associated antigen, by contacting said sample with said cancer associated antigen.



80. Method for screening for a cancerous condition comprising assaying a sample taken from a subject for an immunoreactive cell specific for a peptide derived from a cancer associated antigen encoded by SEQ ID NO: 1, 2, 3, 4, 8, 15, 19, 22 or 26, complexed to an MHC molecule, presence of said immunoreactive cell being indicative of said cancerous condition.

81. An isolated nucleic acid molecule consisting of a nucleotide sequence defined by SEQ ID NO: 1, 2, 3, 8, 15, 19, 22 or 26.

82. Isolated nucleic acid molecule the complimentary sequence of which hybridizes, under stringent conditions, to the nucleotide sequence set forth in SEQ ID NO: 4, 5, 8, 15, 19, 22 or 26.

83. An isolated polypeptide comprising at least 9 consecutive amino acids set forth in SEQ ID NO: 5, 7, 16, 19, 23, 27, or 30.

84. The isolated polypeptide of claim 83, comprising at least 9 consecutive amino acids set forth in SEQ ID NO: 23 or 30.

85. The isolated polypeptide of claim 84, comprising at least 9 consecutive amino acids of the amino acid sequence set forth in SEQ ID NO: 23.

86. The isolated polypeptide of claim 85, comprising amino acids 102-111, 904-912 or 1262-1270 of SEQ ID NO: 23.

87. An isolated nucleic acid molecule which encodes the amino acid sequence of SEQ ID NO: 30.

88. An isolated nucleic acid molecule which encodes the isolated polypeptide of claim 86.

89. Expression vector comprising the isolated nucleic acid molecule of claim 88, operably linked to a promoter.

<110> Jager, Dirk  
Scanlan, Matthew  
Gure, Ali  
Jager, Elke  
Knuth, Alexander  
Old, Lloyd  
Chen, Yao-tseng

<120> Isolated Nucleic Acid Molecules Encoding Cancer Associated Antigens,  
the Antigens per se, and Uses Thereof

<130> LUD 5615.2PCT

<140>

<141>

<150> US 09/451,739

<151> 2000-06-22

<150> US 09/451,739

<151> 1999-11-30

<160> 30

<210> 1  
<211> 1533  
<212> DNA  
<213> Homo sapiens  
<220>  
<221> CDS  
<222> 235  
<400> 1

```
ggttttccac gttggacaag tgcggctcgg cggccagcgg agcgcgcccc ttcccgtgc 60
ccgctccgct cctctcttct acccagccca gtgggcgagt gggcagcggc ggccgcggcg 120
ctgggccctc tcccgcgggt gtgtgcgcgc tcgtacgcgc ggcccccggc gccagccccg 180
ccgcctgaga gggggcctgc gccgccggcc ggggcgtgcg cccggggagcc accgncaccg 240
cggccccgcg cctcaggcgc tggggctccc gcggacccgg aggcggcgga cgggctcggc 300
agatgtagcc gccgggccga agcaggagcc ggccgggggg cgccgggaga gcgagggtt 360
tgcattttgc agtgcattt tttgagggg gcggagggtg gaggaagtcg gaaagccgcg 420
ccgagtcgcc ggggacctcc ggggtgaacc atgttgagtc ctgccaacgg ggagcagctc 480
cacctggtga actatgtgga ggactacctg gactccatcg agtccctgcc ttctgacttg 540
cagagaaatg tctcgtgat gcgggagatc gacgcgaaat accaagagat cctgaaggag 600
ctagacgagt gctacgagcg cttcagtcgc gagacagacg gggcgcagaa gcggcggatg 660
ctgcactgtg tgcagcgcgc gctgatccgc agccaggagc tgggcgacga gaagatccag 720
atcgtgagcc agatggtgga gctggtggag aaccgcacgc ggaggtgga cagccacgtg 780
gagctgttcg aggcgcagca ggagctgggc gacacagcgg gcaacagcgg caaggctggc 840
gcggacaggc ccaaaggcga ggccggcagc caggctgaca agccaacag caagcgtca 900
cggcggcagc gcaacaacga gaaccgtgag aacgcgtcca gcaaccacga ccacgacgac 960
ggcgcctcgg gcacacccaa ggagaagaag gccaaagacct ccaagaagaa gaagcgtcc 1020
aaggccaagg cggagcgaga ggcgtcccct gccgacctcc ccatcgacct caacgaaccc 1080
```

acgtactgtc tgtgcaacca ggtctcctat ggggagatga tcggctgcga caacgacgag 1140  
tgcccatcg agtggttcca cttctcgtgc gtggggctca atcataaacc caagggcaag 1200  
tggtactgtc ccaagtgccg gggggagAAC gagaagacca tggacaaagc cctggagaaa 1260  
tccaaaaaag agagggctta caacaggtag tttgtggaca ggcgcctggt gtgaggagga 1320  
caaaataaac cgtgtattta ttacattgct gcctttgttg aggtgcaagg agtgtaaaat 1380  
gtatattttt aaagaatggt agaaaaggaa ccattccttt catagggatg gcagtgattc 1440  
tgtttgcctt ttgttttcat tggtagacgt gtaacaagaa agtggctctgt ggatcagcat 1500  
tttagaaact acaaatatag gtttgattca aca 1533

&lt;210&gt; 2

&lt;211&gt; 1143

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 2

gagtaaccgc ataatatgcc gttgtccggc acggcgacga gaattcccag atatagcagt 60  
agcagtgatc ccgggacctgt ggctcggggc cggggctgca gttcggaccg cctcccgcga 120  
cccgcggggg ctcggagaca gtttcaggcc gcatccttgc tgacccgagg gtggggccgc 180  
gcgtggccgt ggaaacagat cctgaaggag ctagacgagt gctacgagcg cttcagtcgc 240  
gagacagacg gggcgagaa gcggcgagtg ctgactgtg tgcagcgcg cctgatccgc 300  
agccaggagc tgggagcga gaagatccag atcgtgagcc agatggtgga gctggtggag 360  
aaccgcacgc ggcaggtgga cagccacgtg gagctgttcg aggcgcagca ggagctgggc 420  
gacacagtgg gcaacagcgg caaggttggc gcggacaggc ccaatggcga tgcggtagcg 480  
cagtctgaca agcccaacag caagcgctca cggcgagcgc gcaacaacga gaaccgtgag 540  
aacgcgtcca gcaaccacga ccacgacgac ggcgcctcgg gcacacccaa ggagaagaag 600  
gccaaagacct ccaagaagaa gaagcgctcc aaggccaagg cggagcgaga ggcgtcccct 660  
gccgacctcc ccacgaccc caacgaaccc acgtactgtc tgtgcaacca ggtctcctat 720  
ggggagatga tcggctgcga caacgacgag tgcccatcg agtggttcca cttctcgtgc 780  
gtggggctca atcataaacc caagggcaag tggtagctgc ccaagtgccg gggggagAAC 840  
gagaagacca tggacaaagc cctggagaaa tccaaaaaag agagggctta caacaggtag 900  
tttgtggaca ggcgcctggt gtgaggagga caaaataaac cgtgtattta ttacattgct 960  
gcctttgttg aggtgcaagg agtgtaaaat gtatattttt aaagaatggt agaaaaggaa 1020  
ccattccttt catagggatg gcagtgattc tgtttgcctt ttgttttcat tggtagacgt 1080  
gtaacaagaa agtggctctgt ggatcagcat tttagaaact acaaatatag gtttgattca 1140  
aca 1143

&lt;210&gt; 3

&lt;211&gt; 742

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 3

cgccgtccac accccagcgg ccctgacgct gtcccctccg cgaccctcgc ctctggaaaa 60  
agtgcaggc aaggccacgc ccccgagagg gccggcctcg agcccgagc cccaggggcc 120

tgggacgaga tcctgaagga gctagacgag tgctacgagc gtttcagtcg cgagacagac 180  
 ggggcgacaga agcggcggat gctgcactgt gtgcagcgcg cgctgatccg cagccaggag 240  
 ctgggacgacg agaagatcca gatcgtgagc agatggtgg agctggtgga gaaccgcacg 300  
 cggcaggtgg acagccacgt ggagctgttc gaggcgagc aggagctggg cgacacagcg 360  
 ggcaacagcg gcaaggctgg cgcggacagg cccaaaggcg aggcggcagc gcaggctgac 420  
 aagcccaaca gcaagcgctc acggcgagc cgcaacaacg agaaccgtga gaacgcgtcc 480  
 agcaaccacg accacgacga cggcgctcgg ggcacacca aggagaagaa ggccaagacc 540  
 tccaagaaga agaagcgctc caaggccaag gcgagcgag aggcgtcccc tgccgacctc 600  
 cccatcgacc ccaacgaacc cacgtactgt ctgtgcaacc aggtctccta tggggagatg 660  
 atcggctgcg acaacgacga gtgccccatc gagtgggtcc acttctcgtg cgtggggctc 720  
 aatcataaac ccaagggcaa gt 742

&lt;210&gt; 4

&lt;211&gt; 857

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 4

cctccgagaa cgggtgtccat ggcacagggc gggaagagat aaggcctagg gaaggcgccc 60  
 ctccggccta tccacctctt ctggggctcg gcactaggaa gcagcttccc tctcaggccc 120  
 ctttgtctcc aagccgttcc aaactgagta ccgggagacg acacaaaggg agggcgggtga 180  
 cggtatggcg aggcgcggga gccgcctagg ctgctgggag tgggtggtccg gccgcgggat 240  
 ggagatcctg aaggagctag acgagtgtta cgagcgcttc agtcgcgaga cagacggggc 300  
 gcagaagcgg cggtatgtgc actgtgtgca gcgcgcgtg atccgcagcc aggagctggg 360  
 cgacgagaag atccagatcg tgagccagat ggtggagctg gtggagaacc gcacgcggca 420  
 ggtggacagc cacgtggagc tggtcgagcg gcagcaggag ctgggcgaca cagcgggcaa 480  
 cagcggcaag gctggcgcg agaggcccaa aggcgagggc gcagcgagc ctgacaagcc 540  
 caacagcaag cgctcacggc ggcagcgcaa caacgagaac cgtgagaacg cgtccagcaa 600  
 ccacgaccac gacgacggcg cctcgggcac acccaaggag aagaaggcca agacctccaa 660  
 gaagaagaag cgctccaagg ccaaggcgga gcgagaggcg tcccctgccg acctccccat 720  
 cgaccccaac gaaccacgt actgtctgtg caaccaggtc tcctatgggg agatgatcgg 780  
 ctgcgacaac gacgagtgcc ccacgagtg gttccacttc tcgtgcgtgg ggctcaatca 840  
 taaaccaag ggcaagt 857

&lt;210&gt; 5

&lt;211&gt; 279

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 5

Met Leu Ser Pro Ala Asn Gly Glu Gln Leu His Leu Val Asn Tyr Val  
 1 5 10 15  
 Glu Asp Tyr Leu Asp Ser Ile Glu Ser Leu Pro Phe Asp Leu Gln Arg  
 20 25 30  
 Asn Val Ser Leu Met Arg Glu Ile Asp Ala Lys Tyr Gln Glu Ile Leu  
 35 40 45

Lys Glu Leu Asp Glu Cys Tyr Glu Arg Phe Ser Arg Glu Thr Asp Gly  
 50 55 60  
 Ala Gln Lys Arg Arg Met Leu His Cys Val Gln Arg Ala Leu Ile Arg  
 65 70 75 80  
 Ser Gln Glu Leu Gly Asp Glu Lys Ile Gln Ile Val Ser Gln Met Val  
 85 90 95  
 Glu Leu Val Glu Asn Arg Thr Arg Gln Val Asp Ser His Val Glu Leu  
 100 105 110  
 Phe Glu Ala Gln Gln Glu Leu Gly Asp Thr Val Gly Asn Ser Gly Lys  
 115 120 125  
 Val Gly Ala Asp Arg Pro Asn Gly Asp Ala Val Ala Gln Ser Asp Lys  
 130 135 140  
 Pro Asn Ser Lys Arg Ser Arg Arg Gln Arg Asn Asn Glu Asn Arg Glu  
 145 150 155 160  
 Asn Ala Ser Ser Asn His Asp His Asp Asp Gly Ala Ser Gly Thr Pro  
 165 170 175  
 Lys Glu Lys Lys Ala Lys Thr Ser Lys Lys Lys Lys Arg Ser Lys Ala  
 180 185 190  
 Lys Ala Glu Arg Glu Ala Ser Pro Ala Asp Leu Pro Ile Asp Pro Asn  
 195 200 205  
 Glu Pro Thr Tyr Cys Leu Cys Asn Gln Val Ser Tyr Gly Glu Met Ile  
 210 215 220  
 Gly Cys Asp Asn Asp Glu Cys Pro Ile Glu Trp Phe His Phe Ser Cys  
 225 230 235 240  
 Val Gly Leu Asn His Lys Pro Lys Gly Lys Trp Tyr Cys Pro Lys Cys  
 245 250 255  
 Arg Gly Glu Asn Glu Lys Thr Met Asp Lys Ala Leu Glu Lys Ser Lys  
 260 265 270  
 Lys Glu Arg Ala Tyr Asn Arg  
 275

&lt;210&gt; 6

&lt;211&gt; 210

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 6

Met Leu His Cys Val Gln Arg Ala Leu Ile Arg Ser Gln Glu Leu Gly  
 1 5 10 15  
 Asp Glu Lys Ile Gln Ile Val Ser Gln Met Val Glu Leu Val Glu Asn  
 20 25 30  
 Arg Thr Arg Gln Val Asp Ser His Val Glu Leu Phe Glu Ala Gln Gln  
 35 40 45  
 Glu Leu Gly Asp Thr Val Gly Asn Ser Gly Lys Val Gly Ala Asp Arg  
 50 55 60  
 Pro Asn Gly Asp Ala Val Ala Gln Ser Asp Lys Pro Asn Ser Lys Arg  
 65 70 75 80  
 Ser Arg Arg Gln Arg Asn Asn Glu Asn Arg Glu Asn Ala Ser Ser Asn  
 85 90 95  
 His Asp His Asp Asp Gly Ala Ser Gly Thr Pro Lys Glu Lys Lys Ala  
 100 105 110

Lys Thr Ser Lys Lys Lys Lys Arg Ser Lys Ala Lys Ala Glu Arg Glu  
 115 120 125  
 Ala Ser Pro Ala Asp Leu Pro Ile Asp Pro Asn Glu Pro Thr Tyr Cys  
 130 135 140  
 Leu Cys Asn Gln Val Ser Tyr Gly Glu Met Ile Gly Cys Asp Asn Asp  
 145 150 155 160  
 Glu Cys Pro Ile Glu Trp Phe His Phe Ser Cys Val Gly Leu Asn His  
 165 170 175  
 Lys Pro Lys Gly Lys Trp Tyr Cys Pro Lys Cys Arg Gly Glu Asn Glu  
 180 185 190  
 Lys Thr Met Asp Lys Ala Leu Glu Lys Ser Lys Lys Glu Arg Ala Tyr  
 195 200 205  
 Asn Arg  
 210

<210> 7  
 <211> 235  
 <212> PRT  
 <213> Homo sapiens  
 <400> 7

Met Glu Ile Leu Lys Glu Leu Asp Glu Cys Tyr Glu Arg Phe Ser Arg  
 1 5 10 15  
 Glu Thr Asp Gly Ala Gln Lys Arg Arg Met Leu His Cys Val Gln Arg  
 20 25 30  
 Ala Leu Ile Arg Ser Gln Glu Leu Gly Asp Glu Lys Ile Gln Ile Val  
 35 40 45  
 Ser Gln Met Val Glu Leu Val Glu Asn Arg Thr Arg Gln Val Asp Ser  
 50 55 60  
 His Val Glu Leu Phe Glu Ala Gln Gln Glu Leu Gly Asp Thr Val Gly  
 65 70 75 80  
 Asn Ser Gly Lys Val Gly Ala Asp Arg Pro Asn Gly Asp Ala Val Ala  
 85 90 95  
 Gln Ser Asp Lys Pro Asn Ser Lys Arg Ser Arg Arg Gln Arg Asn Asn  
 100 105 110  
 Glu Asn Arg Glu Asn Ala Ser Ser Asn His Asp His Asp Asp Gly Ala  
 115 120 125  
 Ser Gly Thr Pro Lys Glu Lys Lys Ala Lys Thr Ser Lys Lys Lys Lys  
 130 135 140  
 Arg Ser Lys Ala Lys Ala Glu Arg Glu Ala Ser Pro Ala Asp Leu Pro  
 145 150 155 160  
 Ile Asp Pro Asn Glu Pro Thr Tyr Cys Leu Cys Asn Gln Val Ser Tyr  
 165 170 175  
 Gly Glu Met Ile Gly Cys Asp Asn Asp Glu Cys Pro Ile Glu Trp Phe  
 180 185 190  
 His Phe Ser Cys Val Gly Leu Asn His Lys Pro Lys Gly Lys Trp Tyr  
 195 200 205  
 Cys Pro Lys Cys Arg Gly Glu Asn Glu Lys Thr Met Asp Lys Ala Leu  
 210 215 220  
 Glu Lys Ser Lys Lys Glu Arg Ala Tyr Asn Arg  
 225 230 235

<210> 8  
<211> 772  
<212> DNA  
<213> Homo sapiens  
<221> CDS  
<222> 695,714  
<400> 8  
aaagcgttct cggcggcagc gcaacaacta gaaccgtgag aacgcgtcca gcaaccgcga 60  
cccacgacga cgtcacctcg ggcacgcca aggagaagaa agcccagacc tctaagaaga 120  
agcagggctc catggccaag gcgtagcggc aggcgtcccc cgcagacctc cccatcgacc 180  
ccagcgagcc ctctactgg gagatgatcc gctgcgacaa cgaatgcccc atcgagtggg 240  
tccgcttctc gtgtgtgagt ctcaaccata aaccaaagcg caagtgggtac tgttcagat 300  
gccggggaaa gaacgatggg caaagccctt gagaagtcca gaaaaaaac agggcttata 360  
acaggtagtt tggggacatg cgtctaatag tgaggagaac aaaataagcc agtgtgttga 420  
ttacattgcc acctttgctg aggtgcagga agtgtaaaat gtatatTTTT aaagaatgtt 480  
gttagaggcc gggcgcggtg gctcacgcct gtaatcccag cactttggga ggccgaggcg 540  
gtcggatcac gaggtcagga gatcgagacc atcctggcta acacggtgaa accccgtctc 600  
tactaaaaat tcaaaaaaaa aattagctgg gcgtgggtggc gggcgccctgt agtcccagct 660  
attcgggagg ctgaggcagg agaatggcnt gaacctggga ggtggagctt gcantgagcc 720  
aaggtcgcgc cactgcactc cagcctgggc gacagagcga gactccatct ta 772

<210> 9  
<211> 32  
<212> DNA  
<213> Homo sapiens  
<400> 9  
cacacaggat ccattgttgg tcttgccaac gg 32

<210> 10  
<211> 23  
<212> DNA  
<213> Homo sapiens  
<400> 10  
cgtggtcgtg gttgctggac gcg 23

<210> 11  
<211> 21  
<212> DNA  
<213> Homo sapiens  
<400> 11  
cccagcggcc ctgacgctgt c 21

<210> 12  
<211> 23  
<212> DNA  
<213> Homo sapiens  
<400> 12  
cgtggtcgtg gttgctggac gcg 23

<210> 13  
<211> 23  
<212> DNA  
<213> Homo sapiens  
<400> 13

ggaagagata aggcctaggg aag 23

<210> 14

<211> 23

<212> DNA

<213> Homo sapiens

<400> 14

cggtggtcgtg gttgctggac gcg 23

<210> 15

<211> 2030

<212> DNA

<213> Homo sapiens

<221> CDS

<222> 1628, 1752, 1758, 1769, 1789, 1873, 1908, 1915, 1933, 1970, 1976, 2022

<400> 15

ctcgtgccgt taaagatggt cttctgaagg ctaactgcgg aatgaaagtt tctattccaa 60

ctaaagcctt agaattgatg gacatgcaaa ctttcaaagc agagcctccc gagaagccat 120

ctgccttcga gcctgccatt gaaatgcaaa agtctgttcc aaataaagcc ttggaattga 180

agaatgaaca aacattgaga gcagatgaga tactcccatc agaatccaaa caaaaggact 240

atgaagaaag ttcttgggat tctgagagtc tctgtgagac tgtttcacag aaggatgtgt 300

gtttacccaa ggctacacat caaaaagaaa tagataaaat aaatggaaaa ttagaagagt 360

ctcctgataa tgatgggttt ctgaaggctc cctgcagaat gaaagtttct attccaacta 420

aagccttaga attgatggac atgcaaaact tcaaagcaga gcctcccgag aagccatctg 480

ccttcgagcc tgccattgaa atgcaaaaagt ctgttccaaa taaagccttg gaattgaaga 540

atgaacaaac attgagagca gatcagatgt tcccttcaga atcaaaacaa aagaagggtg 600

aagaaaattc ttgggattct gagagtctcc gtgagactgt ttcacagaag gatgtgtgtg 660

taccaaggc tacacatcaa aaagaaatgg ataaaataag tggaaaatta gaagattcaa 720

ctagcctatc aaaaatcttg gatacagttc attcttgtga aagagcaagg gaacttcaaa 780

aagatcactg tgaacaacgt acaggaaaaa tggaaacaaat gaaaaagaag ttttgtgtac 840

tgaaaaagaa actgtcagaa gcaaaagaaa taaatcaca gttagagaac caaaaagtta 900

aatgggaaca agagctctgc agtgtgagat tgactttaaa ccaagaagaa gagaagagaa 960

gaaatgccga tatattaaat gaaaaaatta gggaagaatt aggaagaatc gaagagcagc 1020

ataggaaaga gttagaagtg aaacaacaac ttgaacaggc tctcagaata caagatatag 1080

aattgaagag tgtagaaagt aatttgaaatc aggtttctca cactcatgaa aatgaaaatt 1140

atctcttaca tgaaaattgc atgttgaaaa aggaaattgc catgctaaaa ctggaaatag 1200

ccacactgaa acaccaatac caggaaaagg aaaataaata ctttgaggac attaagattt 1260

taaaagaaaa gaatgctgaa cttcagatga ccctaaaact gaaagaggaa tcattaacta 1320

aaagggcatc tcaatatagt gggcagctta aagtctgat agctgagaac acaatgctca 1380

cttctaaatt gaaggaaaaa caagacaaag aaataactaga ggcagaaatt gaatcacacc 1440

atcctagact ggcttctgct gtacaagacc atgatcaaat tgtgacatca agaaaaagtc 1500

aagaacctgc tttccacatt gcaggagatg cttgtttgca aagaaaaatg aatgttgatg 1560

tgagtagtac cgatatataa caatgagggt ctccatcaac cactttctga agctcaaagg 1620



aaatccanaa gcctaaaaat taatctcaat tatgcaggag atgctctaag agaaaataca 1680  
 ttggtttcag gaacatgcac aaagagacca acgtgaaaca cagtgtcaaa tgaaggaagc 1740  
 tgaacacatg tntcaaanng aacaagatna tgtgaacaaa cacactganc agcaggagtc 1800  
 tctagatcag aaattatttc aactacaaag caaaaatatg tggcttcaac agcaattagt 1860  
 tcatgcacat aangaaagct gacaacaaaa gcaagataac aattgatntt cattntcttg 1920  
 agaggaaaaat gcncatcatc ttctaaaaga gaaaaatgag gagatatttn attacnataa 1980  
 ccatttaaaa aaccgtata tttcaatatg gaaaaaaaaa anaaaaaaaaa 2030

<210> 16

<211> 513

<212> PRT

<213> Homo sapiens

<400> 16

Met Lys Val Ser Ile Pro Thr Lys Ala Leu Glu Leu Met Asp Met Gln  
1 5 10 15

Thr Phe Lys Ala Glu Pro Pro Glu Lys Pro Ser Ala Phe Glu Pro Ala  
20 25 30

Ile Glu Met Gln Lys Ser Val Pro Asn Lys Ala Leu Glu Leu Lys Asn  
35 40 45

Glu Gln Thr Leu Arg Ala Asp Glu Ile Leu Pro Ser Glu Ser Lys Gln  
50 55 60

Lys Asp Tyr Glu Glu Ser Ser Trp Asp Ser Glu Ser Leu Cys Glu Thr  
65 70 75 80

Val Ser Gln Lys Asp Val Cys Leu Pro Lys Ala Thr His Gln Lys Glu  
85 90 95

Ile Asp Lys Ile Asn Gly Lys Leu Glu Glu Ser Pro Asp Asn Asp Gly  
100 105 110

Phe Leu Lys Ala Pro Cys Arg Met Lys Val Ser Ile Pro Thr Lys Ala  
115 120 125

Leu Glu Leu Met Asp Met Gln Thr Phe Lys Ala Glu Pro Pro Glu Lys  
130 135 140

Pro Ser Ala Phe Glu Pro Ala Ile Glu Met Gln Lys Ser Val Pro Asn  
145 150 155 160

Lys Ala Leu Glu Leu Lys Asn Glu Gln Thr Leu Arg Ala Asp Gln Met  
165 170 175

Phe Pro Ser Glu Ser Lys Gln Lys Lys Val Glu Glu Asn Ser Trp Asp  
180 185 190

Ser Glu Ser Leu Arg Glu Thr Val Ser Gln Lys Asp Val Cys Val Pro  
195 200 205

Lys Ala Thr His Gln Lys Glu Met Asp Lys Ile Ser Gly Lys Leu Glu  
210 215 220

Asp Ser Thr Ser Leu Ser Lys Ile Leu Asp Thr Val His Ser Cys Glu  
225 230 235 240

Arg Ala Arg Glu Leu Gln Lys Asp His Cys Glu Gln Arg Thr Gly Lys  
245 250 255

Met Glu Gln Met Lys Lys Lys Phe Cys Val Leu Lys Lys Lys Leu Ser  
260 265 270

Glu Ala Lys Glu Ile Lys Ser Gln Leu Glu Asn Gln Lys Val Lys Trp

275	280	285
Glu Gln Glu Leu Cys Ser Val Arg Leu Thr Leu Asn Gln Glu Glu Glu 290 295 300		
Lys Arg Arg Asn Ala Asp Ile Leu Asn Glu Lys Ile Arg Glu Glu Leu 305 310 315 320		
Gly Arg Ile Glu Glu Gln His Arg Lys Glu Leu Glu Val Lys Gln Gln 325 330 335		
Leu Glu Gln Ala Leu Arg Ile Gln Asp Ile Glu Leu Lys Ser Val Glu 340 345 350		
Ser Asn Leu Asn Gln Val Ser His Thr His Glu Asn Glu Asn Tyr Leu 325 360 365		
Leu His Glu Asn Cys Met Leu Lys Lys Glu Ile Ala Met Leu Lys Leu 370 375 380		
Glu Ile Ala Thr Leu Lys His Gln Tyr Gln Glu Lys Glu Asn Lys Tyr 385 390 395 400		
Phe Glu Asp Ile Lys Ile Leu Lys Glu Lys Asn Ala Glu Leu Gln Met 405 410 415		
Thr Leu Lys Leu Lys Glu Glu Ser Leu Thr Lys Arg Ala Ser Gln Tyr 420 425 430		
Ser Gly Gln Leu Lys Val Leu Ile Ala Glu Asn Thr Met Leu Thr Ser 435 440 445		
Lys Leu Lys Glu Lys Gln Asp Lys Glu Ile Leu Glu Ala Glu Ile Glu 450 455 460		
Ser His His Pro Arg Leu Ala Ser Ala Val Gln Asp His Asp Gln Ile 465 470 475 480		
Val Thr Ser Arg Lys Ser Gln Glu Pro Ala Phe His Ile Ala Gly Asp 485 490 495		
Ala Cys Leu Gln Arg Lys Met Asn Val Asp Val Ser Ser Thr Asp Ile 500 505 510		

<210> 17  
 <211> 33  
 <212> DNA  
 <213> Homo sapiens  
 <400> 17  
 cacacaggat ccattgcaggc cccgcacaag gag 33

<210> 18  
 <211> 34  
 <212> DNA  
 <213> Homo sapiens  
 <400> 18  
 cacacaaagc ttctaggatt tggcacagcc agag 34

<210> 19  
 <211> 294  
 <212> PRT  
 <213> Homo sapiens  
 <400> 19  
 Met Pro Leu Cys Thr Ala Thr Arg Ile Pro Arg Tyr Ser Ser Ser Ser  
 1 5 10 15  
 Asp Pro Gly Pro Val Ala Arg Gly Arg Gly Cys Ser Ser Asp Arg Leu  
 20 25 30

Pro Arg Pro Ala Gly Pro Ala Arg Arg Gln Phe Gln Ala Ala Ser Leu  
           35                          40                          45  
 Leu Thr Arg Gly Trp Gly Arg Ala Trp Pro Trp Lys Gln Ile Leu Lys  
           50                          55                          60  
 Glu Leu Asp Glu Cys Tyr Glu Arg Phe Ser Arg Glu Thr Asp Gly Ala  
           65                          70                          75                          80  
 Gln Lys Arg Arg Met Leu His Cys Val Gln Arg Ala Leu Ile Arg Ser  
                           85                          90                          95  
 Gln Glu Leu Gly Asp Glu Lys Ile Gln Ile Val Ser Gln Met Val Glu  
                           100                          105                          110  
 Leu Val Glu Asn Arg Thr Arg Gln Val Asp Ser His Val Glu Leu Phe  
           115                          120                          125  
 Glu Ala Gln Gln Glu Leu Gly Asp Thr Val Gly Asn Ser Gly Lys Val  
           130                          135                          140  
 Gly Ala Asp Arg Pro Asn Gly Asp Ala Val Ala Gln Ser Asp Lys Pro  
           145                          150                          155                          160  
 Asn Ser Lys Arg Ser Arg Arg Gln Arg Asn Asn Glu Asn Arg Glu Asn  
                           165                          170                          175  
 Ala Ser Ser Asn His Asp His Asp Asp Gly Ala Ser Gly Thr Pro Lys  
                           180                          185                          190  
 Glu Lys Lys Ala Lys Thr Ser Lys Lys Lys Lys Arg Ser Lys Ala Lys  
           195                          200                          205  
 Ala Glu Arg Glu Ala Ser Pro Ala Asp Leu Pro Ile Asp Pro Asn Glu  
           210                          215                          220  
 Pro Thr Tyr Cys Leu Cys Asn Gln Val Ser Tyr Gly Glu Met Ile Gly  
           225                          230                          235                          240  
 Cys Asp Asn Asp Glu Cys Pro Ile Glu Trp Phe His Phe Ser Cys Val  
                           245                          250                          255  
 Gly Leu Asn His Lys Pro Lys Gly Lys Trp Tyr Cys Pro Lys Cys Arg  
                           260                          265                          270  
 Gly Glu Asn Glu Lys Thr Met Asp Lys Ala Leu Glu Lys Ser Lys Lys  
           275                          280                          285  
 Glu Arg Ala Tyr Asn Arg  
           290                          294

<210> 20  
 <211> 22  
 <212> DNA  
 <213> Homo sapiens  
 <400> 20  
 caaagcagag cctcccgaga ag 22

<210> 21  
 <211> 24  
 <212> DNA  
 <213> Homo sapiens  
 <400> 21  
 cctatgctgc tcttcgattc ttcc 24

<210> 22  
 <211> 4115  
 <212> DNA  
 <213> Homo sapiens  
 <400> 22

ctagtctata cagcaacgac cctacatcgt cactctgggg tcttagaaaag tccataaagc 60  
 tgcctcccg gacaagtcg aagctggaga gatgacaaag ggaagaagac atcaacctta 120  
 atatacaaga gccagaaga gactgctcta actgggcctg gtcaatggcc tgaggaagta 180  
 gtaacatttc ggtagacaga agtgccagct gacgtccttg tggcgaacac ggacacctct 240  
 gatgaaggct tacaatgcca caggaggctt tgcaaatatt tgatagattc ggtgccgata 300  
 taaatctcgt gatgtgtatg caacatggct tccattatgc gtttatagtg gattttgtca 360  
 gtggtggcaa actgctgtcc atggtgcagt atcgaagtgc caacaaggct gcctcacacc 420  
 acttttacta ccataacgaa agaagtgagc aattgtggaa ttttgctgat aaaaatgcaa 480  
 atgcgaatgc gttaataagt taaatgcaca ccctcatgct gctgtatgtc tggatcatca 540  
 gagatagttg catgcttctt agcaaaatgt gacgtctttg tgcatagata gtggagtaac 600  
 tgcagaacat atgctgttac tgtggatttc tcacattcat aacaaattat gaatatatac 660  
 gaaaattatc aaaaatcatc aaataccaat cagaaggaa tctgcaggaa acctgatgag 720  
 gctgcaccct ggcggaaaga cacctgacac gctgaaagct ggtggaaaaa cacctgatga 780  
 ggctgcaccc tgggtggaaag acacctgaca ggctgaaagc tgggtggaaaa acacctgatg 840  
 aggtgcacac ttggtggagg aacatctgac aaattcaatg ttggagaaag gacatctgga 900  
 aagttcgaac gtcagcagaa aaacacctag gaaattacga tcctgcaaaa aaacatctga 960  
 gaaatttacg ggccagcaaa ggaagaccta gaagatcgca gggagaaaaa gaagacacac 1020  
 ctagggaat atgagtcccg aaaagaaaca ctgagaaatt acgtgggcag aaaaggaaga 1080  
 cctaggaaga cgcatgggag aaaaagaaac cctgtaaaga tggatgcgtg caagagtaac 1140  
 atctaataaa cttaaagttt gaaaaaggaa atctaagatg ttgcatgtcc aaaaaagaat 1200  
 catctacaaa gcaagtgcc tgcacagagg tcccatcaga tccaacaag ggaagatgaa 1260  
 gaatattctt tgattctcgg gtctctttga agttctgcaa gattcaagtg gtatacctga 1320  
 gtctatatat aaaaagtaat gagataaata agaagtagaa agcctcctaa aagccatctg 1380  
 ccttcaagcc gccattgaaa gcaaaactct ttccaaataa gcctttgaat gaagaatgaa 1440  
 caaacattga agcagatccg tgttcccacc gaatccaaac aaaggactat aagaaaattc 1500  
 ttgggattct agagtctctg gagactgttt acagaaggat tgtgtttacc aaggctacac 1560  
 atcaaaaaga atagataaaa aaatggaaaa tagaagagtc cctaataaag tggctctctg 1620  
 aaggctacct cggaatgaaa tttctattcc actaaagcct agaattgaag acatgcaaac 1680  
 tttcaaagcg agcctccggg aagccatctg cttcgagcct ccactgaaat caaaagtctg 1740  
 tcccaataa gccttggaat gaaaaatgaa aaacatggag gcagatgaga actcccatca 1800  
 gaatccaaac aaaggactat aagaaaattc tgggatactg gagtctctgt agactgtttc 1860  
 acagaaggat tgtgtttacc aaggctgcgc tcaaaaagaa tagataaaat aatggaaaat 1920  
 tagaagggtc cctgttaaag tggctctctg aggctaactg ggaatgaaag ttctattcca 1980  
 actaaagcct agaattgatg acatgcaaac ttcaaagcag gcctcccgag agccatctgc 2040  
 cttcgagcct ccattgaaat caaagtctg tccaaataa ccttgaatt aagaatgaac 2100  
 aaacattgag gcagatgaga actcccatca aatccaaaca aaggactatg agaaagttct 2160

tggtgattctg gagtctctgt agactgtttc cagaaggatg gtgtttaccc aggctacaca 2220  
tcaaaaagaa tagataaaat aatggaaaat agaagagtct ctgataatga ggttttctga 2280  
aggctccctg agaatgaaag ttctattcga ctaaagcctt gaattgatgg catgcaaact 2340  
ttcaaagcag gcctcccgag agccatctgc ttcgagcctg cattgaaatg aaaagtctgt 2400  
tccaaataaa ccttggaatt aagaatgaac aacattgaga cagatcagat ttcccttcag 2460  
aatcaaaaca aagaagggtg agaaaattct gggattctga agtctccgtg gactgtttca 2520  
cagaaggatg gtgtgtaccc aggctacaca caaaaagaaa ggataaaata gtggaaaatt 2580  
agaagattca ctagcctatc aaaatcttgg tacagttcat cttgtgaaag gcaagggaac 2640  
ttcaaaaaga cactgtgaac acgtacagga aaatggaaca atgaaaaaga gttttgtgta 2700  
ctgaaaaaga actgtcagaa caaaagaaat aaatcacagt agagaacca aagttaaatg 2760  
ggaacaagag tctgcagtgt agattgactt aaaccaagaa aagagaagag agaaatgccg 2820  
atatattaaa gaaaaaatta ggaagaatta gaagaatcga gagcagcata gaaagagtta 2880  
gaagtgaac acaacttgaa aggctctcag atacaagata agaattgaag gtgtagaaag 2940  
taatttgaat aggtttctca actcatgaaa tgaaaattat tcttcatga aattgcatgt 3000  
tgaaaaagga attgccatgc aaaactggaa tagccacact aaacaccaat ccaggaaaag 3060  
gaaaataaat ctttgaggac ttaagatttt aaagaaaaga tgctgaactt agatgaccct 3120  
aaaactgaaa aggaatcatt actaaaagg atctcaatat gtgggcagct aaagttctga 3180  
tagctgagaa acaatgctca ttctaaattg aggaaaaaca gacaaagaaa actagaggca 3240  
gaaattgaat acaccatcct gactggcttc gctgtacaag ccatgatcaa ttgtgacatc 3300  
aagaaaaagt aagaacctgc ttccacattg aggagatgct gtttgcaaag aaaatgaatg 3360  
ttgatgtgag agtacgatat taacaatgag tgctccatca ccactttctg agctcaaagg 3420  
aatccaaaa cctaaaaatt atctcaatta gcaggagatg tctaagagaa atacattgggt 3480  
ttcagaacat cacaaagaga caacgtgaaa acagtgtcaa tgaaggaagc gaacacatgt 3540  
atcaaacga caagataatg gaacaaacac ctgaacagca gagtctctag tcagaaatta 3600  
tttcaactac aagcaaaaat tgtggcttca cagcaattag tcatgcacat agaaagctga 3660  
caacaaaagc agataacaat gatattcatt tcttgagagg aaatgcaaca catctcctaa 3720  
aagagaaaaa gaggagatat taattacaat accatttaaa aaccgtatat tcaatatgaa 3780  
aaagagaaag agaaacagaa actcatgaga acaagcagta gaaacttctt tggagaaaca 3840  
acagaccaga ctttactcac actcatgcta gaggccagtc tagcatcacc tatgttgaaa 3900  
atcttaccac agtctgtgtc acagaatact attttagaag aaaattcatg tttcttctg 3960  
aagcctacag cataaaataa agtg<sup>g</sup>gaaga ttacttggtc cgaattgcat aagctgcaca 4020  
ggattcccat taccctgatg tgcagcagac tcattcaatc aaccagaatc cgctctgcac 4080  
tccagcctag tgacagagtg gactccacct ggaaa 4115

&lt;210&gt; 23

&lt;211&gt; 1341

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 23

Met Thr Lys Arg Lys Lys Thr Ile Asn Leu Asn Ile Gln Asp Ala Gln

1	5	10	15
Lys Arg Thr	Ala Leu His Trp	Ala Cys Val Asn Gly	His Glu Glu Val
	20	25	30
Val Thr Phe	Leu Val Asp Arg	Lys Cys Gln Leu Asp	Val Leu Asp Gly
	35	40	45
Glu His Arg	Thr Pro Leu Met	Lys Ala Leu Gln Cys	His Gln Glu Ala
	50	55	60
Cys Ala Asn	Ile Leu Ile Asp	Ser Gly Ala Asp	Ile Asn Leu Val Asp
	65	70	75
Val Tyr Gly	Asn Met Ala Leu	His Tyr Ala Val	Tyr Ser Glu Ile Leu
	85	90	95
Ser Val Val	Ala Lys Leu Leu	Ser His Gly Ala	Val Ile Glu Val His
	100	105	110
Asn Lys Ala	Ser Leu Thr Pro	Leu Leu Leu Ser	Ile Thr Lys Arg Ser
	115	120	125
Glu Gln Ile	Val Glu Phe Leu	Leu Ile Lys Asn	Ala Asn Ala Asn Ala
	130	135	140
Val Asn Lys	Tyr Lys Cys Thr	Ala Leu Met Leu	Ala Val Cys His Gly
	145	150	155
Ser Ser Glu	Ile Val Gly Met	Leu Leu Gln Gln	Asn Val Asp Val Phe
	165	170	175
Ala Ala Asp	Ile Cys Gly Val	Thr Ala Glu His	Tyr Ala Val Thr Cys
	180	185	190
Gly Phe His	His Ile His Glu	Gln Ile Met Glu	Tyr Ile Arg Lys Leu
	195	200	205
Ser Lys Asn	His Gln Asn Thr	Asn Pro Glu Gly	Thr Ser Ala Gly Thr
	210	215	220
Pro Asp Glu	Ala Ala Pro Leu	Ala Glu Arg Thr	Pro Asp Thr Ala Glu
	225	230	235
Ser Leu Val	Glu Lys Thr Pro	Asp Glu Ala Ala	Pro Leu Val Glu Arg
	245	250	255
Thr Pro Asp	Thr Ala Glu Ser	Leu Val Glu Lys	Thr Pro Asp Glu Ala
	260	265	270
Ala Ser Leu	Val Glu Gly Thr	Ser Asp Lys Ile	Gln Cys Leu Glu Lys
	275	280	285
Ala Thr Ser	Gly Lys Phe Glu	Gln Ser Ala Glu	Glu Thr Pro Arg Glu
	290	295	300
Ile Thr Ser	Pro Ala Lys Glu	Thr Ser Glu Lys	Phe Thr Trp Pro Ala
	305	310	315
Lys Gly Arg	Pro Arg Lys Ile	Ala Trp Glu Lys	Lys Glu Asp Thr Pro
	325	330	335
Arg Glu Ile	Met Ser Pro Ala	Lys Glu Thr Ser	Glu Lys Phe Thr Trp
	340	345	350
Ala Ala Lys	Gly Arg Pro Arg	Lys Ile Ala Trp	Glu Lys Lys Glu Thr
	355	360	365
Pro Val Lys	Thr Gly Cys Val	Ala Arg Val Thr	Ser Asn Lys Thr Lys
	370	375	380
Val Leu Glu	Lys Gly Arg Ser	Lys Met Ile Ala	Cys Pro Thr Lys Glu

385		390		395		400
Ser Ser Thr Lys Ala	Ser Ala Asn Asp Gln Arg Phe Pro Ser Glu Ser					
	405		410			415
Lys Gln Glu Glu Asp	Glu Glu Tyr Ser Cys Asp Ser Arg Ser Leu Phe					
	420		425			430
Glu Ser Ser Ala Lys Ile	Gln Val Cys Ile Pro Glu Ser Ile Tyr Gln					
	435		440			445
Lys Val Met Glu Ile Asn Arg	Glu Val Glu Glu Pro Pro Lys Lys Pro					
	450		455			460
Ser Ala Phe Lys Pro Ala Ile	Glu Met Gln Asn Ser Val Pro Asn Lys					
	465		470			480
Ala Phe Glu Leu Lys Asn Glu	Gln Thr Leu Arg Ala Asp Pro Met Phe					
	485		490			495
Pro Pro Glu Ser Lys Gln Lys	Asp Tyr Glu Glu Asn Ser Trp Asp Ser					
	500		505			510
Glu Ser Leu Cys Glu Thr Val	Ser Gln Lys Asp Val Cys Leu Pro Lys					
	515		520			525
Ala Thr His Gln Lys Glu Ile	Asp Lys Ile Asn Gly Lys Leu Glu Glu					
	530		535			540
Ser Pro Asn Lys Asp Gly Leu Leu Lys	Ala Thr Cys Gly Met Lys Val					
	545		550			560
Ser Ile Pro Thr Lys Ala Leu Glu Leu Lys	Asp Met Gln Thr Phe Lys					
	565		570			575
Ala Glu Pro Pro Gly Lys Pro Ser Ala Phe	Glu Pro Ala Thr Glu Met					
	580		585			590
Gln Lys Ser Val Pro Asn Lys Ala Leu Glu Leu Lys	Asn Glu Gln Thr					
	595		600			605
Trp Arg Ala Asp Glu Ile Leu Pro Ser Glu Ser Lys	Gln Lys Asp Tyr					
	610		615			620
Glu Glu Asn Ser Trp Asp Thr Glu Ser Leu Cys Glu Thr Val Ser	Gln					
	625		630			640
Lys Asp Val Cys Leu Pro Lys Ala Ala His Gln Lys Glu Ile Asp Lys						
	645		650			655
Ile Asn Gly Lys Leu Glu Gly Ser Pro Val Lys Asp Gly Leu Leu Lys						
	660		665			670
Ala Asn Cys Gly Met Lys Val Ser Ile Pro Thr Lys Ala Leu Glu Leu						
	675		680			685
Met Asp Met Gln Thr Phe Lys Ala Glu Pro Pro Glu Lys Pro Ser Ala						
	690		695			700
Phe Glu Pro Ala Ile Glu Met Gln Lys Ser Val Pro Asn Lys Ala Leu						
	705		710			720
Glu Leu Lys Asn Glu Gln Thr Leu Arg Ala Asp Glu Ile Leu Pro Ser						
	725		730			735
Glu Ser Lys Gln Lys Asp Tyr Glu Glu Ser Ser Trp Asp Ser Glu Ser						
	740		745			750
Leu Cys Glu Thr Val Ser Gln Lys Asp Val Cys Leu Pro Lys Ala Thr						
	755		760			765
His Gln Lys Glu Ile Asp Lys Ile Asn Gly Lys Leu Glu Glu Ser Pro						

770					775					780					
Asp 785	Asn	Asp	Gly	Phe	Leu 790	Lys	Ala	Pro	Cys	Arg 795	Met	Lys	Val	Ser	Ile 800
Pro	Thr	Lys	Ala	Leu 805	Glu	Leu	Met	Asp	Met 810	Gln	Thr	Phe	Lys	Ala	Glu 815
Pro	Pro	Glu	Lys 820	Pro	Ser	Ala	Phe	Glu 825	Pro	Ala	Ile	Glu	Met	Gln	Lys
Ser	Val	Pro 835	Asn	Lys	Ala	Leu	Glu 840	Leu	Lys	Asn	Glu	Gln	Thr	Leu	Arg
Ala	Asp 850	Gln	Met	Phe	Pro	Ser 855	Glu	Ser	Lys	Gln	Lys 860	Lys	Val	Glu	Glu
Asn 865	Ser	Trp	Asp	Ser	Glu 870	Ser	Leu	Arg	Glu	Thr 875	Val	Ser	Gln	Lys	Asp 880
Val	Cys	Val	Pro	Lys 885	Ala	Thr	His	Gln	Lys 890	Glu	Met	Asp	Lys	Ile	Ser 895
Gly	Lys	Leu 900	Glu	Asp	Ser	Thr	Ser	Leu 905	Ser	Lys	Ile	Leu	Asp	Thr	Val
His	Ser	Cys 915	Glu	Arg	Ala	Arg	Glu 920	Leu	Gln	Lys	Asp	His	Cys	Glu	Gln
Arg	Thr 930	Gly	Lys	Met	Glu	Gln 935	Met	Lys	Lys	Lys	Phe 940	Cys	Val	Leu	Lys
Lys 945	Lys	Leu	Ser	Glu	Ala 950	Lys	Glu	Ile	Lys	Ser 955	Gln	Leu	Glu	Asn	Gln 960
Lys	Val	Lys	Trp	Glu 965	Gln	Glu	Leu	Cys	Ser 970	Val	Arg	Leu	Thr	Leu	Asn 975
Gln	Glu	Glu	Glu	Lys 980	Arg	Arg	Asn	Ala 985	Asp	Ile	Leu	Asn	Glu	Lys	Ile 990
Arg	Glu	Glu 995	Leu	Gly	Arg	Ile	Glu 1000	Glu	Gln	His	Arg	Lys	Glu	Leu	Glu 1005
Val	Lys 1010	Gln	Gln	Leu	Glu	Gln 1015	Ala	Leu	Arg	Ile	Gln	Asp	Ile	Glu	Leu 1020
Lys 1025	Ser	Val	Glu	Ser	Asn 1030	Leu	Asn	Gln	Val	Ser 1035	His	Thr	His	Glu	Asn 1040
Glu	Asn	Tyr	Leu	Leu 1045	His	Glu	Asn	Cys	Met	Leu	Lys	Lys	Glu	Ile	Ala 1055
Met	Leu	Lys	Leu	Glu 1060	Ile	Ala	Thr	Leu	Lys 1065	His	Gln	Tyr	Gln	Glu	Lys 1070
Glu	Asn	Lys 1075	Tyr	Phe	Glu	Asp	Ile 1080	Lys	Ile	Leu	Lys	Glu	Lys	Asn	Ala 1085
Glu	Leu	Gln	Met	Thr	Leu	Lys 1095	Leu	Lys	Glu	Glu	Ser	Leu	Thr	Lys	Arg 1100
Ala	Ser	Gln	Tyr	Ser	Gly 1110	Gln	Leu	Lys	Val	Leu	Ile	Ala	Glu	Asn	Thr 1120
Met	Leu	Thr	Ser	Lys 1125	Leu	Lys	Glu	Lys	Gln	Asp	Lys	Glu	Ile	Leu	Glu 1135
Ala	Glu	Ile	Glu	Ser	His 1140	His	Pro	Arg	Leu	Ala	Ser	Ala	Val	Gln	Asp 1150
His	Asp	Gln	Ile	Val	Thr	Ser	Arg	Lys	Ser	Gln	Glu	Pro	Ala	Phe	His



1155	1160	1165
Ile Ala Gly Asp Ala Cys Leu Gln Arg Lys Met Asn Val Asp Val Ser 1170	1175	1180
Ser Thr Ile Tyr Asn Asn Glu Val Leu His Gln Pro Leu Ser Glu Ala 1185	1190	1200
Gln Arg Lys Ser Lys Ser Leu Lys Ile Asn Leu Asn Tyr Ala Gly Asp 1205	1210	1215
Ala Leu Arg Glu Asn Thr Leu Val Ser Glu His Ala Gln Arg Asp Gln 1220	1225	1230
Arg Glu Thr Gln Cys Gln Met Lys Glu Ala Glu His Met Tyr Gln Asn 1235	1240	1245
Glu Gln Asp Asn Val Asn Lys His Thr Glu Gln Gln Glu Ser Leu Asp 1250	1255	1260
Gln Lys Leu Phe Gln Leu Gln Ser Lys Asn Met Trp Leu Gln Gln Gln 1265	1270	1280
Leu Val His Ala His Lys Lys Ala Asp Asn Lys Ser Lys Ile Thr Ile 1285	1290	1295
Asp Ile His Phe Leu Glu Arg Lys Met Gln His His Leu Leu Lys Glu 1300	1305	1310
Lys Asn Glu Glu Ile Phe Asn Tyr Asn Asn His Leu Lys Asn Arg Ile 1315	1320	1325
Tyr Gln Tyr Glu Lys Glu Lys Ala Glu Thr Glu Asn Ser 1330	1335	1340

<210> 24  
 <211> 22  
 <212> DNA  
 <213> Homo sapiens  
 <400> 24  
 aatgggaaca agagctctgc ag 22

<210> 25  
 <211> 23  
 <212> DNA  
 <213> Homo sapiens  
 <400> 25  
 gggtcatctg aagttcagca ttc 23

<210> 26  
 <211> 3673  
 <212> DNA  
 <213> Homo sapiens  
 <221> CDS  
 <222> 439, 473, 1789  
 <400> 26  
 caagagcttg gcgatacaga aatttctgct ggtgttgggg cgggtgcggg aactgaagac 60  
 gggcgagtgc gagccggggg cgggtgctgg ggaagggtaa gcgggaagcg agggcgaggg 120  
 gtaggggctg gggaaagggcg agcgggagcg gcgggctctc tctagcaggg ggctgcagcc 180  
 atgaagaggc tcttagctgc cgctggcaag ggcgtgcggg gcccggagcc cccgaacccc 240  
 ttcagcgaac ggggtctacac tgagaaggac tacgggacca tctacttcgg ggatctaggg 300  
 aagatccata cagctgcctc ccggggccaa gtccagaagc tggagaagat gacagtaggg 360  
 aagaagcccg tcaacctgaa caaagagat atgaagaaga ggactgctct acactggggc 420

tgtgtcaatg gccatgcana agtagtaaca tttctggtag acagaaagtg ccngcttaat 480  
gtccttgatg gcgaaggag gacacctctg atgaaggctc tacaatgcga gagggaagct 540  
ttgtgcaaatt attctcatag atgctggtgc tgatctaaat tatgtagatg tgtatggcaa 600  
cacggctctc cattatgccg tttatagtga gaatttatta atggtggcaa cactgctgtc 660  
ctatggtgca gtcacgagg tgcaaaacaa ggctagcctc acaccccttt tactggccat 720  
acagaaaaga agcaagcaaa ctgtggaatt tttactaaca aaaaatgcaa atgcaaacgc 780  
atttaatgag tctaaatgca cagccctcat gcttgccata tgtgaaggct catcagagat 840  
agtcggcatg cttcttcagc aaaatgttga cgtctttgct gaagacatac atggaataac 900  
tgagaacgt tatgctgctg ctogtggagt taattacatt catcaacaac ttttggaaca 960  
tatacgaaaa ttacctaaaa atcctcaaaa taccatcca gaaggaaat ctacaggaa 1020  
acctgatgag gctgcaccct tggcggaag aacacctgac acggctgaaa gcttgctgga 1080  
aaaaacacct gacgaggctg cacgcttggg ggagggaacg tctgccaaa ttcaatgtct 1140  
ggggaagca acatctggaa agtttgaaca gtcaacagaa gaaacaccta ggaaaatttt 1200  
gaggctaca aaagaacat ctgagaaatt tcatggcca gcaaaagaaa gatctaggaa 1260  
gatcacatgg gagaaaaag aaacatctgt aaagactgaa tgcgtggcag gagtaacacc 1320  
taataaaact gaagttttg aaaaggaac atctaatatg attgcatgtc ctacaaaaga 1380  
aacatctaca aaagcaagta caaatgtgga tgtgagttct gtagagccta tattcagtct 1440  
ttttggcaca cggactattg aaaattcaca gtgtacaaa gttgaggaag actttaatct 1500  
tgctaccaag attatctota agagtgtgc acagaattat acgtgtttac ctgatgtac 1560  
atatcaaaaa gatatcaaaa caataaatca caaaatagaa gatcagatgt tcccatcaga 1620  
atccaaacga gaggaagatg aagaatattc ttgggattct gggagtctct ttgagagttc 1680  
tgcaagact caagtgtgta tacctgagtc tatgtatcag aaagtaatgg agataaatag 1740  
agaagtagaa gagcttcctg agaagccatc tgccttcaag cctgccgtng aaatgcaaaa 1800  
gactgttcca aataagcct ttgaattgaa gaatgaacaa acattgagag cagctcagat 1860  
gttcccatca gaatccaaac aaaaggacga tgaagaaaat tcttgggatt ctgagagtcc 1920  
ctgtgagacg gtttcacaga aggatgtgta tttacccaaa gctacacatc aaaaagaatt 1980  
cgatacctta agtggaatat tagaagagtc tcctgttaaa gatggtcttc tgaagcctac 2040  
ctgtggaagg aaagtttctc ttccaaataa agccttagaa ttaaaggaca gagaaacatt 2100  
caaagcagag tctcctgata aagatggtct tctgaagcct acctgtggaa ggaaagtctc 2160  
tcttccaaat aaagccttag aattaaagga cagagaaaca ctcaaagcag agtctcctga 2220  
taatgatggt cttctgaagc ctacctgtgg aaggaaagtt tctcttccaa ataaagcttt 2280  
agaattgaag gacagagaaa cattcaaagc agctcagatg ttcccatcag aatccaaaca 2340  
aaaggatgat gaagaaaatt cttgggattt tgagagtctt cttgagactc tcttacagaa 2400  
tgatgtgtgt ttacccaagg ctacacatca aaaagaattc gataccttaa gtggaaaatt 2460  
agaagagtct cctgataaag atggtcttct gaagcctacc tgtggaatga aaatttctct 2520  
tccaaataaa gccttagaat tgaaggacag agaaacattc aaagcagagg atgtgagttc 2580

tgtagagtcc acattcagtc tttttggcaa accgactact gaaaattcac agtctacaaa 2640  
 agttgaggaa gactttaatc ttactaccaa ggaggagca acaaagacag taactggaca 2700  
 acaggaacgt gatattggca ttattgaacg agtccacaa gatcaacaa ataagatgcc 2760  
 cacatcagaa ttaggaagaa aagaagatac aaaatcaact tcagattctg agattatctc 2820  
 tgtgagtgat acacagaatt atgagtgttt acctgaggct acatatcaaa aagaaataaa 2880  
 gacaacaaat ggcaaaatag aagagtctcc tgaaaagcct tctcactttg agcctgccac 2940  
 tgaaatgcaa aactctgttc caaataaagg cttagaatgg aagaataaac aaacattgag 3000  
 agcagattca actaccctat caaaaatctt ggatgcactt cttcttctgt aaagagggaag 3060  
 ggaacttaaa aaagataact gtgaacaaat tacagcaaaa atggaacaaa tgaaaaataa 3120  
 gttttgtgta ctacaaaagg aactgtcaga agcgaagaa ataaaatcac agttagagaa 3180  
 ccaaaaagct aaatgggaac aagagctctg cagtgtgaga ttgcctttaa atcaagaaga 3240  
 agagaagaga agaaatgtcg atatattaaa agaaaaaatt agaccogaag agcaacttag 3300  
 gaaaaagtta gaagtgaac accaacttga acagactctc agaatacaag atatagaatt 3360  
 gaaaagtgt acaagtaatt tgaatcaggt ttctcacact catgaaagtg aaaatgatct 3420  
 ctttcatgaa aattgcatgt tgaaaaagga aattgccatg ctaaaactgg aagtagccac 3480  
 actgaaacat caacaccagg tgaaggaaaa taaatacttt gaggacatta agattttaca 3540  
 agaaaagaat gctgaacttc aaatgaccct aaaactgaaa cagaaaacag taacaaaaag 3600  
 ggcatctcag tatagagagc agcttaaagt tctgacggca gagaacacga tgctgacttc 3660  
 taaattgaag gaa 3673

&lt;210&gt; 27

&lt;211&gt; 1011

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 27

Met Val Ala Thr Leu Leu Ser Tyr Gly Ala Val Ile Glu Val Gln Asn  
 1 5 10 15  
 Lys Ala Ser Leu Thr Pro Leu Leu Leu Ala Ile Gln Lys Arg Ser Lys  
 20 25 30  
 Gln Thr Val Glu Phe Leu Leu Thr Lys Asn Ala Asn Ala Asn Ala Phe  
 35 40 45  
 Asn Glu Ser Lys Cys Thr Ala Leu Met Leu Ala Ile Cys Glu Gly Ser  
 50 55 60  
 Ser Glu Ile Val Gly Met Leu Leu Gln Gln Asn Val Asp Val Phe Ala  
 65 70 75 80  
 Glu Asp Ile His Gly Ile Thr Ala Glu Arg Tyr Ala Ala Ala Arg Gly  
 85 90 95  
 Val Asn Tyr Ile His Gln Gln Leu Leu Glu His Ile Arg Lys Leu Pro  
 100 105 110  
 Lys Asn Pro Gln Asn Thr Asn Pro Glu Gly Thr Ser Thr Gly Thr Pro  
 115 120 125  
 Asp Glu Ala Ala Pro Leu Ala Glu Arg Thr Pro Asp Thr Ala Glu Ser  
 130 135 140  
 Leu Leu Glu Lys Thr Pro Asp Glu Ala Ala Arg Leu Val Glu Gly Thr

145		150		155		160
Ser Ala Lys Ile	Gln Cys Leu Gly Lys	Ala Thr Ser Gly Lys	Phe Glu			
	165	170	175			
Gln Ser Thr	Glu Thr Pro Arg Lys	Ile Leu Arg Pro Thr	Lys Glu			
	180	185	190			
Thr Ser Glu Lys	Phe Ser Trp Pro Ala Lys	Glu Arg Ser Arg Lys	Ile			
	195	200	205			
Thr Trp Glu Glu	Lys Glu Thr Ser Val Lys	Thr Glu Cys Val Ala Gly				
	210	215	220			
Val Thr Pro Asn Lys	Thr Glu Val Leu Glu Lys	Gly Thr Ser Asn Met				
	225	230	235			240
Ile Ala Cys Pro	Thr Lys Glu Thr Ser Thr	Lys Ala Ser Thr Asn Val				
	245	250	255			
Asp Val Ser	Ser Val Glu Pro Ile Phe	Ser Leu Phe Gly Thr Arg Thr				
	260	265	270			
Ile Glu Asn Ser	Gln Cys Thr Lys Val Glu Glu	Asp Phe Asn Leu Ala				
	275	280	285			
Thr Lys Ile Ile	Ser Lys Ser Ala Ala Gln	Asn Tyr Thr Cys Leu Pro				
	290	295	300			
Asp Ala Thr Tyr	Gln Lys Asp Ile Lys Thr	Ile Asn His Lys Ile Glu				
	305	310	315			320
Asp Gln Met Phe	Pro Ser Glu Ser Lys Arg	Glu Glu Asp Glu Glu Tyr				
	325	330	335			
Ser Trp Asp Ser	Gly Ser Leu Phe Glu Ser	Ser Ala Lys Thr Gln Val				
	340	345	350			
Cys Ile Pro Glu	Ser Met Tyr Gln Lys Val Met	Glu Ile Asn Arg Glu				
	355	360	365			
Val Glu Glu Leu	Pro Glu Lys Pro Ser Ala Phe	Lys Pro Ala Val Glu				
	370	375	380			
Met Gln Lys Thr	Val Pro Asn Lys Ala Phe	Glu Leu Lys Asn Glu Gln				
	385	390	395			400
Thr Leu Arg Ala	Ala Gln Met Phe Pro Ser	Glu Ser Lys Gln Lys Asp				
	405	410	415			
Asp Glu Glu Asn	Ser Trp Asp Ser Glu Ser Pro	Cys Glu Thr Val Ser				
	420	425	430			
Gln Lys Asp Val	Tyr Leu Pro Lys Ala Thr His	Gln Lys Glu Phe Asp				
	435	440	445			
Thr Leu Ser Gly	Lys Leu Glu Glu Ser Pro Val	Lys Asp Gly Leu Leu				
	450	455	460			
Lys Pro Thr Cys	Gly Arg Lys Val Ser Leu Pro	Asn Lys Ala Leu Glu				
	465	470	475			480
Leu Lys Asp Arg	Glu Thr Phe Lys Ala Glu Ser	Pro Asp Lys Asp Gly				
	485	490	495			
Leu Leu Lys Pro	Thr Cys Gly Arg Lys Val Ser	Leu Pro Asn Lys Ala				
	500	505	510			
Leu Glu Leu Lys	Asp Arg Glu Thr Leu Lys Ala	Glu Ser Pro Asp Asn				
	515	520	525			
Asp Gly Leu Leu	Lys Pro Thr Cys Gly Arg Lys	Val Ser Leu Pro Asn				

530	535	540
Lys Ala Leu Glu Leu	Lys Asp Arg Glu Thr	Phe Lys Ala Ala Gln Met
545	550	555 560
Phe Pro Ser Glu Ser	Lys Gln Lys Asp Asp	Glu Glu Asn Ser Trp Asp
	565	570 575
Phe Glu Ser Phe Leu Glu Thr	Leu Leu Gln Asn Asp	Val Cys Leu Pro
	580 585	590
Lys Ala Thr His Gln Lys Glu Phe Asp Thr	Leu Ser Gly Lys Leu Glu	
	595 600	605
Glu Ser Pro Asp Lys Asp Gly Leu Leu Lys Pro	Thr Cys Gly Met Lys	
	610 615	620
Ile Ser Leu Pro Asn Lys Ala Leu Glu Leu Lys	Asp Arg Glu Thr Phe	
	625 630	635 640
Lys Ala Glu Asp Val Ser Ser Val Glu Ser Thr	Phe Ser Leu Phe Gly	
	645	650 655
Lys Pro Thr Thr Glu Asn Ser Gln Ser Thr Lys	Val Glu Glu Asp Phe	
	660 665	670
Asn Leu Thr Thr Lys Glu Gly Ala Thr Lys Thr	Val Thr Gly Gln Gln	
	675 680	685
Glu Arg Asp Ile Gly Ile Ile Glu Arg Ala Pro	Gln Asp Gln Thr Asn	
	690 695	700
Lys Met Pro Thr Ser Glu Leu Gly Arg Lys Glu	Asp Thr Lys Ser Thr	
	705 710	715 720
Ser Asp Ser Glu Ile Ile Ser Val Ser Asp Thr	Gln Asn Tyr Glu Cys	
	725	730 735
Leu Pro Glu Ala Thr Tyr Gln Lys Glu Ile Lys	Thr Thr Asn Gly Lys	
	740 745	750
Ile Glu Glu Ser Pro Glu Lys Pro Ser His Phe	Glu Pro Ala Thr Glu	
	755 760	765
Met Gln Asn Ser Val Pro Asn Lys Gly Leu Glu	Trp Lys Asn Lys Gln	
	770 775	780
Thr Leu Arg Ala Asp Ser Thr Thr Leu Ser Lys	Ile Leu Asp Ala Leu	
	785 790	795 800
Pro Ser Cys Glu Arg Gly Arg Glu Leu Lys Lys	Asp Asn Cys Glu Gln	
	805 810	815
Ile Thr Ala Lys Met Glu Gln Met Lys Asn Lys	Phe Cys Val Leu Gln	
	820 825	830
Lys Glu Leu Ser Glu Ala Lys Glu Ile Lys Ser	Gln Leu Glu Asn Gln	
	835 840	845
Lys Ala Lys Trp Glu Gln Glu Leu Cys Ser Val	Arg Leu Pro Leu Asn	
	850 855	860
Gln Glu Glu Glu Lys Arg Arg Asn Val Asp Ile	Leu Lys Glu Lys Ile	
	865 870	875 880
Arg Pro Glu Glu Gln Leu Arg Lys Lys Leu Glu	Val Lys His Gln Leu	
	885 890	895
Glu Gln Thr Leu Arg Ile Gln Asp Ile Glu Leu	Lys Ser Val Thr Ser	
	900 905	910
Asn Leu Asn Gln Val Ser His Thr His Glu Ser	Glu Asn Asp Leu Phe	

915	920	925
His Glu Asn Cys Met Leu Lys Lys Glu Ile Ala Met Leu Lys Leu Glu 930 935 940		
Val Ala Thr Leu Lys His Gln His Gln Val Lys Glu Asn Lys Tyr Phe 945 950 955 960		
Glu Asp Ile Lys Ile Leu Gln Glu Lys Asn Ala Glu Leu Gln Met Thr 965 970 975		
Leu Lys Leu Lys Gln Lys Thr Val Thr Lys Arg Ala Ser Gln Tyr Arg 980 985 990		
Glu Gln Leu Lys Val Leu Thr Ala Glu Asn Thr Met Leu Thr Ser Lys 995 1000 1005		
Leu Lys Glu 1010		

<210> 28  
 <211> 23  
 <212> DNA  
 <213> Homo sapiens  
 <400> 28  
 tctcatagat gctggtgctg atc 23

<210> 29  
 <211> 24  
 <212> DNA  
 <213> Homo sapiens  
 <400> 29  
 cccagacatt gaattttggc agac 24

<210> 30  
 <211> 56  
 <212> PRT  
 <213> Homo sapiens  
 <400> 30

Met Glu Glu Ile Ser Ala Ala Ala Val Lys Val Val Pro Gly Pro Glu 1 5 10 15
Arg Pro Ser Pro Phe Ser Gln Leu Val Tyr Thr Ser Asn Asp Ser Tyr 20 25 30
Ile Val His Ser Gly Asp Leu Arg Lys Ile His Lys Ala Ala Ser Arg 35 40 45
Gly Gln Val Arg Lys Leu Glu Lys 50 55